

RESEARCH SUMMARY

David C. Johnston
March 2004

The current primary focus of our current work is to synthesize new and/or improved strongly correlated electron metallic oxide compounds and carry out physical property measurements on them to search for novel physical behaviors and/or ground states, and to clarify the origins of interesting properties of known materials. In the following some background in our research area and some of our progress is outlined. Literature references by other groups have the prefix “L” and are listed at the end of this narrative. Our papers in this resume are referenced as either straight numerals for our papers in the list of refereed publications or preceded by “O” for our “Other Publications”. All three lists of references are given below.

High T_c Cuprates

The discovery of superconducting transition temperatures T_c up to about 35 K in what turned out to be the layered cuprate $\text{La}_{2-x}\text{Ba}_x\text{CuO}_4$ by Bednorz and Müller in 1986 [L1-L3] stimulated extraordinary worldwide experimental and theoretical research activity to understand the physics of the materials and generate new compounds. The physical properties of this and related families of materials were found to be highly anisotropic, reflecting the layered crystal structures in which the Cu atoms form a square lattice as shown in the left panel of Fig. 1 where an oxygen atom resides between each nearest-neighbor pair of copper atoms. Our discovery in early 1987 of long-range antiferromagnetic order at about 250 K in the undoped insulating parent compound La_2CuO_4 [92, 94, 95] demonstrated that the Cu atoms carry local magnetic moments as expected from an ionic model for the d^9 spin-1/2 Cu^{+2} ions. Our observation of a continuous evolution in the temperature dependent magnetic susceptibility with doping level x in the similar superconducting $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ system [118] and inelastic magnetic neutron scattering investigations on this system [L4] indicated that the Cu local moments survive the insulator to superconducting metal transition with increasing x . The undoped parent compounds of subsequently studied families of layered cuprate superconductors were also found to become antiferromagnetic, as illustrated by the discoveries of antiferromagnetism by the PI and collaborators in $\text{YBa}_2\text{Cu}_3\text{O}_6$ [99, 106], $\text{Ca}_{0.85}\text{Sr}_{0.15}\text{CuO}_2$ [122] and $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ [128]. (For an extensive review of the magnetic properties of the single-layer cuprates, see [O23]). These observations indicated that the cuprate high T_c superconductors should be viewed as doped antiferromagnetic insulators. This situation was unprecedented in the history of superconductivity, where superconductivity was previously precluded by the presence of a dense array of local magnetic moments on the same sublattice of the structure on which the superconductivity would have resided. This history in turn suggested that a different mechanism from the conventional electron-phonon interaction may be responsible for the high T_c superconductivity observed in the cuprates. A leading candidate, following the above observations, has been an electronic mechanism involving Cu spin interactions. However, there is still no consensus on the mechanism, so it is difficult to predict where to look for the next high T_c superconductor system or family. The main thrust of the present work is to attempt to synthesize new oxide materials in the search for novel behaviors and ground states, including high temperature superconductivity. The ultimate goal in the latter context is to synthesize a superconducting system with a transition temperature above room temperature, necessarily leading to new physics and hopefully allowing superconductivity to become beneficial and ubiquitous in power transmission, electronic devices, and generally in our everyday lives.




Fig. 1 (color) (Left figure) Comparison between the body-centered tetragonal K_2NiF_4 ($I4/mmm$) crystal structure of La_2CuO_4 (above 530 K, called the T structure), the orthorhombic ($Immm$) crystal structure of Sr_2CuO_3 , and the body-centered tetragonal ($I4/mmm$) crystal structure of Nd_2CuO_4 (called the T' structure).

Fig. 2 (Right figure) Structural, magnetic and superconducting phase diagram of the $La_{2-x}Sr_xCuO_4$ system [O23].

In addition to substitutional doping of the insulating cuprate parent compounds as in the $La_{2-x}(Ca,Sr,Ba)_xCuO_4$ systems, anionic doping, usually involving oxygen addition or intercalation, has been utilized to produce superconducting samples. The most important example is the $YBa_2Cu_3O_{6+x}$ ($0 \leq x \leq 1$) system noted above where hole-doping is achieved for $x > 0$ and superconductivity is observed for x larger than about 0.4 [O12, O13]. The optimum T_c of around 93 K is achieved for x close to 1. In a related vein, we discovered new phases that were induced by hydrogen reduction and by subsequent oxidation of Ln_2CuO_4 compounds ($Ln = La, Pr, Nd, Sm, Eu, Gd$) [132]. Another important example of an oxygen-doped system is the La_2CuO_{4+x} ($0 \leq x \leq 0.12$) system where T_c up to about 45 K is attained for the larger x values. We have synthesized large homogeneous polycrystalline [150] and single crystal [158] samples using room-temperature electrochemical oxidation of La_2CuO_4 in aqueous base. In a collaboration with J. D. Jorgensen's group at Argonne National Laboratory who carried out structural studies using powder neutron diffraction, we found that the excess oxygen occupies empty sites between the CuO_2 layers [153] which are in fact completely occupied in the related Nd_2CuO_4 structure shown in the right-hand panel of Fig. 1. We also mapped out the structural (T, x) phase diagram of this system using the polycrystalline samples [162]. A miscibility gap was discovered between $x = 0.011$ and $x = 0.055$; antiferromagnetic ordering occurs below the low end and superconductivity above the high end compositions of the miscibility gap, with a two-phase mixture in between, which was confirmed by subsequent single crystal neutron diffraction studies [160]. This phase diagram is distinctly different from that of the $La_{2-x}Sr_xCuO_4$ system (Fig. 2; for a review, see [O23]), which has no such miscibility gap versus doping level x . We have also determined the temperature dependence of the oxygen diffusion rate in La_2CuO_{4+x} ; oxygen diffusion was found to be significant to remarkably low temperatures of around 200 K [192]. R. J. Birgeneau's group at MIT later carried out neutron diffraction structural studies on one of our superconducting La_2CuO_{4+x} crystals and on other crystals synthesized there [190]. They discovered that for compositions beyond

the miscibility gap, the intercalated oxygen is “staged”, i.e., occupies the space only between every n^{th} CuO_2 layer ($n = 4, 6, \dots$), where the stage number n depends on the excess oxygen concentration x .

An important and interesting aspect of the hole doped high T_c cuprates is that the doped holes have often been found to occur in a non-random distribution within the CuO_2 planes. Some of the earliest evidence for this was found in 1992-1993 from our observed scaling of the magnetic susceptibility versus temperature and doping level x of the lightly-doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ system, where the doped holes were inferred to segregate into thin one-dimensional walls separating undoped, locally antiferromagnetic domains [151]. Additional evidence for this configuration was found from ^{139}La NQR measurements carried out in collaboration with F. Borsa’s group [146, 154, 182]. Neutron diffraction evidence for the existence of static doped-hole segregation was obtained by a Brookhaven group in 1995 on the compound $\text{La}_{1.48}\text{Nd}_{0.4}\text{Sr}_{0.12}\text{CuO}_4$ [L5]. The one-dimensional doped-hole configurations separating undoped locally antiferromagnetic domains are now called “stripes”. “Stripe physics” has become ubiquitous in the high T_c field (for a review, see [O23]). It is not yet clear whether the occurrence of stripes helps or hinders the high temperature superconductivity in the layered cuprate systems.

As is well-known, the tetragonal K_2NiF_4 (T) structure compound La_2CuO_4 (left panel in Fig. 1) can be hole-doped but not electron-doped, whereas the tetragonal Nd_2CuO_4 (T’) structure compounds like Nd_2CuO_4 (right panel in Fig. 1) can be electron-doped but not hole-doped (see [O23]). Chlorine can sometimes be incorporated into these structures. The T-structure tetragonal Cl-containing compound $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ exists in which the two apical oxygen atoms in La_2CuO_4 are replaced by Cl and the La by Sr, leaving the CuO_2 layers intact. This compound remains tetragonal to low temperatures (instead of distorting to a lower symmetry structure like La_2CuO_4 does below 530 K [92]) and has been important in establishing the intrinsic magnetic properties of the undistorted CuO_2 square lattice [127, 128, 130, 131, 144, 181, 191, 204]. We have carried out experiments attempting to electron-dope T-structure La_2CuO_4 by partially substituting O by Cl to form $\text{La}_2\text{CuO}_{4-y}\text{Cl}_y$ (J. M. Hill and D. C. Johnston, 2003, unpublished). The solid state syntheses were carried out in a variety of ways and at various temperatures using either CuCl or LaOCl as Cl sources. Consistent with all previous work on this compound, we found that electron doping of the T structure did not occur using the described method.

The $\text{La}_{2-x}\text{Ca}_x\text{CuO}_4$ system exhibits superconductivity up to temperatures of about 20 K for $x \geq 0.10$ [L3]. The system was found to be slightly oxygen deficient [L37]. Remarkably, nothing at all is currently known about the normal state magnetic properties or magnetic phase diagram of this hole-doped system. Therefore, we began a study of the magnetic susceptibility and oxygen content of this system to determine if there are any interesting qualitative differences with those of the well-studied $\text{La}_{2-x}(\text{Ba}, \text{Sr})_x\text{CuO}_4$ systems (J. M. Hill and D. C. Johnston, 2003-2004, unpublished). From the orthorhombicity of these orthorhombic samples, defined as $2(b-a)/(a+b)$, versus Ca concentration x , we deduce that the tetragonal-to-orthorhombic structural transition temperature T_o decreases through room temperature (300 K) at a Ca concentration $x \approx 0.17$. By contrast, T_o in the $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ system decreases below room temperature at a much lower Sr concentration $x \approx 0.10$ (see [O23]). The oxygen contents of our samples were determined by reduction under hydrogen in a Perkin-Elmer thermal gravimetric analyzer. There is no indication that the samples are oxygen deficient to within the error of ± 0.04 in the formula $\text{La}_{2-x}\text{Ca}_x\text{CuO}_4$, contrary to the results in [L37]. The magnetic susceptibility, however, is a much more sensitive measure of the oxygen content. La_2CuO_4 samples synthesized in air generally exhibit both a reduced Néel temperature T_N and a tiny volume fraction of superconductivity due to the absorption of excess oxygen from the air during synthesis. This indicates that the amount of excess oxygen in the $\text{La}_2\text{CuO}_{4+x}$ samples is just within the miscibility gap between antiferromagnetic and superconducting compositions for $0.011 \leq x \leq 0.055$ as discussed in the previous section. From magnetic susceptibility measurements, the samples show both antiferromagnetic

ordering at the peaks of the susceptibility and a trace of superconductivity (< 0.1 vol %) below about 40 K evidently due to excess oxygen absorbed from the air during synthesis. Studies of $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ showed a systematic decrease of T_N followed by the disappearance of long-range antiferromagnetic order at $x \approx 0.02$ (see [O23]). A similar smooth systematic decrease with increasing x does not occur for $\text{La}_{2-x}\text{Ca}_x\text{CuO}_4$, perhaps due to differences in the oxygen contents of the various samples. Experiments are in progress to remove the excess oxygen from the whole series of samples by annealing them in an inert gas and to extend our studies to higher Ca concentrations.

Heavy Fermion Spinel Compound LiV_2O_4 and Related Materials

An important question regarding the origin of high temperature superconductivity in the layered cuprates is: what is so special about copper? One of our strategies for researching this question and to search for additional novel physics is to examine systems in which the spin $S = 1/2$ d^9 Cu^{+2} ion is replaced by 3d transition metal cations with the same spin $S = 1/2$ but with a d^1 configuration instead, so that now there is one electron in the 3d shell instead of one hole. An ideal candidate for this purpose is vanadium, which shows stable oxidation states in oxides of +2, +3, +4 (especially), and +5. The cation V^{+4} has $S = 1/2$ with one electron in the d shell. Our initial studies were of the mixed perovskite system LaVO_3 - SrVO_3 . We hoped to see novel physics emerge near the known composition-induced metal-insulator transition, which was not realized [147, 148].

We then moved on to look at the enigmatic compound LiV_2O_4 with the fcc normal spinel structure, where the vanadium cations formally have a $d^{1.5}$ configuration [O17]. Due to the non-integral oxidation state of the V cations and their crystallographic equivalence, symmetry demands that the compound be metallic, which in fact was found long ago from resistivity measurements on small hydrothermally grown single crystals [L6]. Despite the metallic character, the magnetic susceptibility was known to show a Curie-Weiss behavior down to low temperatures (4 K) corresponding to a local magnetic moment system with $S = 1/2$ and spectroscopic splitting factor g close to 2 [L7]. This behavior suggested that one of the 1.5 d -electrons per vanadium was somehow localized on each V atom and the remaining 0.5 d -electron per vanadium was itinerant and responsible for the metallic character. The Weiss temperature in the Curie-Weiss behavior was around -60 K, and yet long range antiferromagnetic ordering was not observed above 4 K. In fact, the V sublattice of the structure consists of corner-sharing tetrahedra, themselves consisting of V triangles, so the V sublattice is strongly geometrically frustrated for antiferromagnetic ordering and the possibility therefore existed that long-range ordering would be completely precluded. If so, how would the spin entropy of the system be dissipated upon cooling? Is new physics involved?

This question led us to an intensive investigation of the synthesis, characterization, and properties of high-purity LiV_2O_4 which resulted in our discovery in 1997 of this material as the first clear example of a d -electron heavy fermion system as reflected in the structural, magnetic susceptibility, specific heat, muon spin relaxation, and NMR properties below about 25 K (the “Kondo” or “coherence” temperature T_K) [205]. Thus the answer to the motivating question posed above is that the spin entropy of the presumed V local moments is converted on cooling below T_K to entropy of the conduction carriers which is manifested as a huge (0.44 J/mol K^2) zero temperature electronic specific heat coefficient as shown in Fig. 3. As the heavy fermion state develops below T_K , the local moment behavior of the V sublattice is transformed to large nearly temperature-independent Pauli paramagnetism as shown in Fig. 4 (the Curie-Weiss behavior below 25 K previously reported by others was due to a contribution from magnetic defects and/or impurities). The heavy fermion behavior of LiV_2O_4 was completely unexpected by the condensed matter physics community since the extended nature of the magnetic V d -orbitals (compared to f -orbitals in f -electron heavy fermion compounds) and their expected strong hybridization with oxygen $2p$ orbitals would seem to preclude such behavior. The presence of potentially new physics was indicated.




Fig. 3 (Left figure) Electronic specific heat C_e divided by temperature T versus T for high purity LiV_2O_4 . The lattice contribution has been subtracted. The extrapolated zero temperature limit of C_e/T is 0.44 J/mol K^2 .

Fig. 4 (Right figure) Magnetic susceptibility χ versus temperature T for high-purity LiV_2O_4 . The weak broad maximum occurs at 16 K. A Curie-Weiss behavior occurs above about 50 K.

Subsequent additional measurements of the above properties and also of the thermal expansion were carried out over the next few years which all self-consistently supported our original heavy fermion interpretation for the low-temperature properties of LiV_2O_4 [207, 210, 216, 217, 224]. In addition, electrical resistivity measurements down to 0.3 K on small hydrothermally grown single crystals by a group in Japan showed a T^2 dependence as expected for a Fermi liquid, with an extremely large coefficient consistent with a very heavy Fermi liquid [L8-L10]. Their specific heat, magnetic susceptibility, ^7Li NMR, and thermal expansion measurements on the crystals [L8-L10] were all consistent with our previous results on polycrystalline samples. They also found that the Hall coefficient shows a distinct peak at about 15 K and crosses over from hole-like to electron-like on heating above 50 K [L8, L10]. Two inelastic magnetic neutron scattering studies have been carried out on polycrystalline samples [L11-L14]. Both studies found behaviors at low temperatures consistent with those known for f -electron heavy fermion compounds. They disagreed, however, on the nature of the magnetic state above T_K . Krimmel et al. [L11-L13] characterized this behavior as that of a nearly *ferromagnetic* metal, whereas the more complete later data of Lee et al. indicated that the heavy fermion state develops on cooling from a state with short-range *antiferromagnetic* spin correlations as expected from the antiferromagnetic Weiss temperature in the Curie-Weiss magnetic susceptibility behavior above T_K [L14]. Inelastic magnetic neutron scattering studies on large single crystals are clearly called for to provide more complete information on the generalized susceptibility and magnetic excitations both below and above T_K . Crystals of the required minimum size (at least 1 g) have not been synthesized up to now. A plethora of theories have been advanced to explain the mechanism for formation of the heavy fermion mass in LiV_2O_4 below T_K , but there is as yet no consensus on the correct approach (for a brief review of some of the theories, see [224]).

Many experiments of various types on high purity polycrystalline and single crystal samples of the metallic transition metal spinel-structure oxide compound LiV_2O_4 showed heavy fermion behaviors from 0.3 K to around 20 K as just described. However, claims have appeared in the literature [L15-L17] that a different behavior occurs in the ^7Li NMR spin-lattice relaxation rate (and other properties) below 1 K from the heavy fermion behavior we observed in this quantity above 1 K. In addition, they

found a non-exponential (“stretched exponential”) decay to the nuclear magnetization in their pulsed NMR experiments, whereas we observed and reported a strictly exponential decay. To address these claims and understand the origin of their divergent results, we extended our ^7Li NMR measurements of this compound down to 0.5 K (S.-H. Baek, F. Borsa, S. Kondo, and D. C. Johnston, unpublished work). We found the temperature dependence of the nuclear spin-lattice relaxation rate to be consistent with our previous results for a high-purity sample with an equivalent $S = 1/2$ magnetic defect concentration of about $n = 0.01$ mol%. In addition, we found that a sample with a higher magnetic defect concentration $n = 0.3$ mol% shows a distinctly different behavior that is similar to that of Refs. [L15-L17]. Finally, we found a strictly exponential decay of the nuclear magnetization for the high purity sample but which becomes non-exponential as the magnetic defect concentration increases. In fact, we had already shown in 1997 from our early μSR studies of two samples with different magnetic defect concentrations that static spin glass ordering occurs below 0.8 K when $n \sim 0.3\%$, whereas no static spin ordering occurs above 0.02 K for our highest purity sample with $n \sim 0.01\%$ [205]. These results indicate that (i) the ground state of high purity LiV_2O_4 is a heavy fermion ground state with heavy fermion behaviors at low temperatures as we originally reported, (ii) the ground state of LiV_2O_4 is qualitatively changed from a heavy fermion ground state to some type of spin-glass state as the concentration of magnetic defects in the material increases above about 0.1 mol%, and (iii) the authors of [L15-L17] carried out their experiments on samples with high magnetic defect concentrations and did not measure the intrinsic properties of pure LiV_2O_4 .

As an extension of our previous work on polycrystalline samples of LiV_2O_4 , in preliminary experiments we attempted to grow single crystals of this compound using fused salt electrolysis of molten electrolytes with composition $(\text{Li}_2\text{O})_x\text{V}_2\text{O}_5$ with $x = 0.5\text{-}1.0$, in 50 mL Pt crucibles under Ar atmosphere at 600-650 °C with currents of 0.1-10 mA. This technique instead consistently yielded single crystals of the known compound LiV_2O_5 , as determined from single crystal x-ray diffraction measurements, with dimensions up to $1 \times 2 \times 15$ mm³. Single crystals of this material had been grown before using flux growth as reported in the literature, but our crystal growth method as applied to this compound is new.

Of the hundreds of known spinel structure transition metal oxide compounds, only two remain metallic to low temperatures: LiV_2O_4 and LiTi_2O_4 . Superconductivity was discovered in 1973 by the PI at temperatures up to 13.7 K in LiTi_2O_4 [5, 12, 13]. Until the advent of high- T_c cuprates in 1986, this was the highest T_c oxide compound. The cation-substituted Ti spinel systems $\text{Li}[\text{Li}_x\text{Ti}_{2-x}]\text{O}_4$ and $\text{Li}[\text{Al}_x\text{Ti}_{2-x}]\text{O}_4$ undergo composition-induced metal-to-insulator transitions as x increases. In collaboration with R. J. Gooding’s group at Queen’s University, we studied these transitions using a quantum site percolation model [230]. The results suggested that strong electronic correlations are both present and important in these Ti oxide spinel systems, just as they obviously are in LiV_2O_4 .

Pyrochlores

We have extended our studies of the fcc spinel compound LiV_2O_4 to include synthesis and properties of pyrochlore structure materials of general composition $\text{A}_2\text{B}_2\text{O}_6\text{O}'$, where A is usually a lanthanide and B a transition metal. The O and O’ atoms occupy crystallographically distinct lattice sites. Both the A and B sublattices are identical to the V sublattice in LiV_2O_4 . However, the A atoms are 8-fold coordinated by O whereas the B atoms have 6-fold octahedral coordination by O. For antiferromagnetically coupled B atoms, the B sublattice is geometrically frustrated for long-range antiferromagnetic ordering, as in LiV_2O_4 . Hypothetical metallic members of this structure class containing local magnetic moments therefore have the potential to exhibit heavy fermion behavior and thus be included in our new class of d -electron heavy fermion compounds. The known pyrochlore compound $\text{Lu}_2\text{V}_2\text{O}_7$ [L18, L19] contains d^1 $S = 1/2$ V^{+4} cations and is a ferromagnetic n-type

semiconductor with a Curie temperature $T_C = 73$ K and an activation energy of 0.2 eV [L19-L23]. We carried out a clean synthesis of hypothetical $\text{Lu}_2\text{V}_2\text{O}_2\text{F}$ (where F replaces O') in sealed molybdenum tubes at 1250°C , which would formally contain $d^{1.5}$ V cations as in LiV_2O_4 and would be metallic if it formed. Unfortunately, the compound did not form (G. Knoke, J. M. Hill, and D. C. Johnston, 2002, unpublished).

We discovered a new series $\text{Lu}_2\text{V}_2\text{O}_{7-x}$ of oxygen deficient pyrochlores with $0.4 \leq x \leq 0.7$ by annealing $\text{Lu}_2\text{V}_2\text{O}_7$ in flowing H_2 (G. Knoke, J. M. Hill, and D. C. Johnston, unpublished work). Magnetization measurements showed that T_C for the oxygen-deficient materials is much lower than the value of 73 K for the $\text{Lu}_2\text{V}_2\text{O}_7$ parent compound, reaching 15 K for $x = 0.7$. Above T_C , the magnetic susceptibility follows the Curie-Weiss law with a Curie constant consistent with a mixture of localized spin $S = 1/2$ V^{+4} and $S = 1$ V^{+3} cations dictated by the oxygen deficiency x in an ionic model. Although we could not carry out electronic transport measurements due to the powder nature of these samples, from the composition dependence of the Curie constant we infer that the series does not become metallic within the composition range investigated. X-ray crystallographic studies demonstrated that the oxygen deficiency is accompanied by Lu-V antisite disorder, which is reflected in a strong reduction in the intensities of the x-ray reflections with (odd odd odd) Miller indices. An outstanding question is why the Curie temperature decreases with increasing oxygen deficiency, even though the average spin of the vanadium sublattice increases with oxygen deficiency. The explanation may have its origins in the antisite disorder just mentioned and deserves further study.

Spin Chain Oxide Compounds

The field of low-dimensional magnetism received a major boost following the discovery of the cuprate high T_c superconductors and of the magnetic character of the parent and doped compounds discussed above. For example, it was discovered theoretically in 1988 that the ground state of the spin $S = 1/2$ square lattice Heisenberg antiferromagnet has long-range antiferromagnetic order (but only at temperature $T = 0$ for an isolated layer) [L23, L24]. A surprising finding from conformal field theory in 1994 was that the magnetic susceptibility of the fiducial $S = 1/2$ Heisenberg chain initially increases with temperature from $T = 0$ with infinite slope, arising from a low-temperature “logarithmic correction” to the susceptibility of the form $1/\log(T^*/T)$, where T^* is a characteristic temperature [L25]. Our numerical calculations support this finding and in addition yield the next few higher order logarithmic correction terms [222, 223]. We also carried out very high accuracy calculations and simulations of the magnetic susceptibility versus temperature of both the uniform antiferromagnetic $S = 1/2$ Heisenberg chain and the alternating exchange chain in which the antiferromagnetic exchange interaction between adjacent spins alternates between two values J_1 and J_2 . We obtained a high-accuracy two-dimensional function describing the susceptibility versus temperature for $0 \leq J_1/J_2 \leq$ infinity by fitting the numerical data to within the accuracy of the data [222]. This high-accuracy function will be of great use in the future for fitting experimental susceptibility data for uniform and alternating-exchange $S = 1/2$ chain compounds (see the fits of the susceptibility of Sr_2CuO_3 below and of $(\text{VO})_2\text{P}_2\text{O}_7$ in the next section). The new high-accuracy numerical results on the magnetic susceptibility versus temperature of the uniform chain [L25, 222, 223] supplant the venerable Bonner-Fisher calculations from 1964 [L26], where the prediction is too low by as much as 10% at the lower temperatures (see Fig. 8.1 in [O23]). For reviews of these and other theoretical developments on the antiferromagnetically coupled $S = 1/2$ spin chain and square lattice, see [O23] and [222].

On the experimental side, in 1995 we discovered that the linear chain compound Sr_2CuO_3 is a $S = 1/2$ Heisenberg chain with a very large Cu-Cu exchange coupling $J \sim 2200$ K [173, 203]. The structure of this compound (middle panel of Fig. 1) is derived from that of the tetragonal high temperature K_2NiF_4 structure of La_2CuO_4 (left panel of Fig. 1) by removing entire lines of O atoms

within the CuO_2 layers along one of the (100) directions and substituting Sr for La, resulting in an orthorhombic unit cell. From our fits to the magnetic susceptibility versus temperature, the nearest-neighbor antiferromagnetic exchange coupling strength between the $S = 1/2$ copper spins is $J = 2150(150)$ K [173, 203]. This J value is about 50% larger than in the above layered cuprate parent compounds (~ 1500 - 1600 K, see [O23]), even though the Cu-O-Cu bonding is nearly identical in the two cases (we address this disparity below). Long-range antiferromagnetic order due to interchain coupling does not set in until the temperature decreases to $T_N = 5.4$ K [L19], so the ratio $T_N/J = 0.002$ is very small. The extrapolated zero-temperature ordered moment from magnetic neutron diffraction measurements on single crystals is only $0.06(3)$ μ_B/Cu [L27]. The expected saturation moment is $\mu_{\text{sat}} = gS\mu_B = 1.1$ μ_B/Cu , and the much lower value observed was attributed to quantum fluctuations associated with the quasi-one-dimensionality of the spin system [L27]. These properties together establish Sr_2CuO_3 as one of the best model $S = 1/2$ quasi-one-dimensional antiferromagnetic Heisenberg spin chain systems to date. Up to now there are no reports of doping this compound into the metallic state. As is well-known, the physics of one-dimensional antiferromagnets is quite different in certain respects from higher-dimensional systems. We are very interested in following the evolution in the physical properties of one-dimensional antiferromagnets such as linear chains and spin ladders (below) with doping to compare and contrast the results with the extensive results on the two-dimensional layered cuprates. We anticipate that the nature of the evolution may be quite different than in higher-dimensional systems and that new physics might be accessible via this route.

We tried to dope the quasi-1D spin-1/2 Heisenberg chain compound Sr_2CuO_3 using electrochemical oxidation in aqueous base solution (J. M. Hill and D. C. Johnston, unpublished work). As described above, we successfully used this technique previously to electrochemically dope holes into insulating quasi-2D La_2CuO_4 to produce superconducting $\text{La}_2\text{CuO}_{4+y}$ ($0.06 \leq y \leq 0.12$) powders and crystals. We found, however, that Sr_2CuO_3 decomposes when exposed to either water vapor or liquid water, the main decomposition product being $\text{Sr}_2\text{Cu}(\text{OH})_6$. The structure of this compound [L28] consists of isolated $\text{Cu}(\text{OH})_6$ octahedra. We carried out a thorough study of the EPR, magnetization versus magnetic field at fixed temperatures T and magnetic susceptibility χ versus T of $\text{Sr}_2\text{Cu}(\text{OH})_6$ [228]. The $\chi(T)$ data are well described by a Curie-Weiss law, with a small Weiss temperature $\theta \approx -3$ K that corresponds to a weak antiferromagnetic Cu-Cu superexchange interaction $J \approx 1$ K. The formation of $\text{Sr}_2\text{Cu}(\text{OH})_6$ on the surfaces of Sr_2CuO_3 samples exposed to air is therefore likely to be the previously unknown source of the variable amounts of Curie-like contributions to the susceptibility of Sr_2CuO_3 previously documented by us [173]. $\text{Sr}_2\text{Cu}(\text{OH})_6$ also serves as a nice reference material for comparison with the magnetic properties of the much more strongly interacting ($J \sim 2000$ K) high- T_c cuprates. We then tried to dope holes into Sr_2CuO_3 via electrochemical oxidation in a nonaqueous electrolyte consisting of tetramethylammonium hydroxide in methyl alcohol, a technique reported to be successful for oxidizing La_2CuO_4 [L29], but no evidence that doping occurred was found from either crystallographic or magnetic susceptibility measurements.

The ambient pressure (AP) phase of the compound vanadyl pyrophosphate, $(\text{VO})_2\text{P}_2\text{O}_7$, contains spin-1/2 d^1 V^{+4} ions. The PI and collaborators discovered in 1987 from magnetic susceptibility data that this compound exhibits a spin gap and modeled the magnetic susceptibility versus temperature data in terms of an antiferromagnetic alternating-exchange spin chain model [90]. For a review of the subsequent convoluted history of the study and interpretation of the magnetic properties of this compound, which for a time was considered to be a spin ladder compound, see [227]. Ultimately our original model that we used to fit the susceptibility data now appears to be the correct one. In collaboration with experimentalists in Japan, we have measured and modeled new magnetic susceptibility versus temperature data for both polycrystalline and single crystal samples of this compound, and also of a new high-pressure (HP) modification of this compound discovered in 1999



Trellis Layer Structure of SrCu_2O_3 and CaV_2O_5 

Fig. 5 (Left figure) The “trellis layer” structure of the Cu and V atoms in SrCu_2O_3 and CaV_2O_5 . Within each two-leg ladder, an oxygen atom is situated between each nearest-neighbor pair of transition metal atoms.

Fig. 6 (Right figure) Measured spin susceptibility of SrCu_2O_3 (dots), together with our fit (solid curve) on the basis of the nearest-neighbor Heisenberg model. Also shown is an ab initio calculation [35] in which a four-spin cyclic exchange interaction and a weak diagonal antiferromagnetic interaction is included in addition to the principal Heisenberg exchange interaction. The exchange interactions for each fit are shown in the figure.

[225], to determine the magnetic exchange interactions between the V spins [225, 227]. To do these fits, the PI used his new high-accuracy two-dimensional function for the susceptibility versus temperature and J_1/J_2 of $S = 1/2$ uniform and alternating-exchange Heisenberg chains discussed above. In a *tour de force* using this function to fit the experimental data, we found that the AP phase contains *two* distinct types of alternating-exchange V^{+4} chains, each with its own spin gap and pair of alternating antiferromagnetic exchange interactions. This result is consistent with the crystal structure, which indicates the presence of two distinguishable spin chains, and with the magnon dispersion relations found from previous inelastic neutron scattering measurements [L30]. In addition, the new HP phase was found to contain a single type of V^{+4} $S = 1/2$ alternating-exchange chain, again consistent with that crystal structure.

Spin Ladder Oxide Compounds

Another offshoot of the high T_c field was spin ladder physics which began in earnest as a subfield of low-dimensional magnetism in 1993 (for reviews, see [O23] and [230A]). An n -leg spin ladder is a spin configuration in which the spins lie on the vertices where the n legs (or rails) meet the perpendicular rungs of the ladder. A square spin lattice, as in the layered high T_c cuprates, is one limiting case with $n = \text{infinity}$, and a linear spin chain is the other limit with $n = 1$. An early theoretical prediction was that $S = 1/2$ ladders with even n should have an energy gap (“spin gap”) between a (nonmagnetic) singlet ground state and the lowest-lying (magnetic) triplet excited states, whereas ladders with odd n should be gapless [e.g., L31, L32]. This prediction was confirmed via magnetic susceptibility and other measurements on the $S = 1/2$ spin ladder compounds SrCu_2O_3 with $n = 2$ and

$\text{Sr}_2\text{Cu}_3\text{O}_5$ with $n = 3$ [L33] (and of course on the above Sr_2CuO_3 with $n = 1$), all of which are insulators. The “trellis layer” structure of SrCu_2O_3 is shown in Fig. 5; the two-leg ladder compound CaV_2O_5 has a similar structure within the V_2O_3 planes. The 180° Cu-O-Cu superexchange path within a ladder is strong and antiferromagnetic, whereas the 90° Cu-O-Cu path between ladders is weakly ferromagnetic and is frustrating, leading to an effective magnetic isolation of the ladders from each other. From the crystal chemistry of these materials, one expects that the Cu-Cu superexchange coupling should be about the same in all three cuprate compounds, with the rung (J') and leg (J) couplings about the same with $J'/J \sim 1$ [L31, L32]. However, in 1996 we discovered that the (rather sparse) existing magnetic susceptibility versus temperature calculations *based on the nearest-neighbor Heisenberg model* for the $S = 1/2$ two-leg ladder, when compared with the experimental magnetic susceptibility data for SrCu_2O_3 , indicated instead that $J'/J \sim 0.5$, with $J \sim 2000$ K [198]. This J value is on the order of that in the linear chain compound Sr_2CuO_3 but larger than the values of 1500-1600 K previously inferred for the square lattice cuprate parent compounds [O23]. However, the *average* of J' and J is about 1500 K. These conundrums were further addressed and confirmed both theoretically and experimentally by the PI and collaborators in very extensive work on spin ladders described next.

We recently completed an extensive five-year combined theoretical and experimental study, in collaboration with scientists in Germany, Switzerland, Russia, and Japan, of the magnetic susceptibility versus temperature $\chi(T)$ of insulating spin $S = 1/2$ two- and three-leg Heisenberg ladders and ladder oxide compounds [230A]. The $\chi(T)$ of isolated two-leg ladders with spatially anisotropic antiferromagnetic (AF) Heisenberg exchange was calculated by quantum Monte Carlo (QMC) simulations, with and without ferromagnetic (FM) second-neighbor diagonal intraladder coupling, and for two-leg ladders coupled into a two-dimensional (2D) stacked ladder configuration or a 3D $\text{LaCuO}_{2.5}$ -type interladder coupling configuration. We obtained accurate analytical fits and interpolations of these data and of previously reported related QMC $\chi(T)$ simulation data for the isolated ladder with spatially isotropic exchange, for the 2D trellis layer configuration and for isotropic and anisotropic three-leg ladders. We also calculated the one- and two-magnon dispersion relations for the isolated 2×12 ladder with $0.5 \leq J'/J \leq 1$, where J is the AF coupling constant along the legs and J' is that along the rungs. The exchange constants in the two-leg ladder compound SrCu_2O_3 were estimated from LDA+U calculations. On the experimental side, we determined the detailed crystal structures of SrCu_2O_3 and of the three-leg ladder compound $\text{Sr}_2\text{Cu}_3\text{O}_5$. New experimental $\chi(T)$ data were obtained for SrCu_2O_3 and $\text{LaCuO}_{2.5}$, and for the nominally two-leg ladder vanadites CaV_2O_5 and MgV_2O_5 which are structurally similar to SrCu_2O_3 . These and literature $\chi(T)$ data for these compounds and for $\text{Sr}_2\text{Cu}_3\text{O}_5$ were modeled using our QMC simulation fits. CaV_2O_5 was found to be essentially a dimer compound with AF intradimer coupling 669(3) K ($g = 1.96$), in agreement with previous results [L34]. The leg and rung exchange constants found for isostructural MgV_2O_5 are very different from those in CaV_2O_5 , as predicted previously from LDA+U calculations. For SrCu_2O_3 , we found that $J'/J = 0.5$ and $J = 1900$ K, assuming a spectroscopic splitting factor $g = 2.1$ (see the fit in Fig. 6), confirming our previous 1996 modeling results [198]. The interladder coupling $J''/J = 0.01(1)$ perpendicular to the ladder layers was found to be very weak and on the spin-gapped side of the quantum critical point (QCP) at $J_{\text{QCP}}''/J = 0.048(2)$ for $J'/J = 0.5$. The three-leg ladder compound $\text{Sr}_2\text{Cu}_3\text{O}_5$ was also found to exhibit strong intraladder exchange anisotropy, with $J'/J = 0.66(5)$ and $J = 1810(150)$ K for $g = 2.1(1)$. The $\chi(T)$ data for $\text{LaCuO}_{2.5}$ were consistent with $J'/J = 0.5$ with $J = 1700$ K, again assuming $g = 2.1$, and with a 3D FM interladder coupling $J^{\text{3D}}/J = -0.05$ which is close to and on the AF ordered side of the QCP at $J_{\text{QCP}}^{\text{3D}}/J = -0.036(1)$ for $J'/J = 0.5$, consistent with the observed AF-ordered ground state.

The surprisingly strong spatial anisotropy of the bilinear exchange constants within the cuprate spin ladders, $J'/J \sim 0.5$, derived unambiguously from our fits of the experimental $\chi(T)$ data by precise predictions of the Heisenberg model as described above, is in conflict with expectation from crystal chemical arguments which predict that $J'/J \geq 1$. The resolution of this conundrum is likely that the Heisenberg Hamiltonian is insufficient to describe the magnetism of the cuprate ladder compounds. In particular, we suggested that a four-spin cyclic exchange interaction is also present which has a strong influence on $\chi(T)$. This hypothesis is supported by calculations of Mizuno, Tohyama, and Maekawa that indicate the presence of this interaction in SrCu_2O_3 [L35]. They found that a relatively small amount (9%) of this interaction relative to J has a profound influence on $\chi(T)$ and that the observed $\chi(T)$ is well reproduced by including both the Heisenberg and cyclic exchange interactions, as shown previously in Fig. 6. One important implication is that the latter interaction is also important to the magnetic properties of the square lattice cuprates. This is consistent with the conclusion of, e.g., Coldea et al. who found that this term must be included in the Hamiltonian in their fits to their inelastic magnetic neutron scattering measurements of the magnon dispersion relations in La_2CuO_4 [L36]. Another important implication is that the exchange interaction J inferred for the square lattice and spin chain cuprates on the basis of the nearest-neighbor Heisenberg model alone is only an *effective* interaction, which is too small relative to the actual value and includes the influence of the cyclic exchange interaction (see the exchange parameters in Fig. 6). This explains why the J in the linear chain cuprate Sr_2CuO_3 (2200 K), in which a cyclic exchange interaction cannot be present, is significantly larger than the (effective) J of 1500-1600 K inferred previously (see [O23]) for the square lattice cuprates.

The Layered Orthogonal Dimer Compound $\text{SrCu}_2(\text{BO}_3)_2$.

We have been carrying out an extensive series of experiments (J. M. Hill, S. Sanger, M. Tillman, S. Madison, and D. C. Johnston, unpublished work) to try to dope the 2D spin dimer compound $\text{SrCu}_2(\text{BO}_3)_2$ [L38]. This insulating compound has a spin gap and is of great current experimental and theoretical interest in its own right because it is the only known physical realization [L39] of the so-called Shastry-Sutherland model [L40] for a particular layered frustrated dimer spin system, for which the magnetic ground state is known exactly. Due to the geometric frustration for antiferromagnetic ordering inherent in the structure (shown in Fig. 7) [L39, L41], this compound has qualitatively different magnetic properties [L38] and interactions [L39, L42] than in the undoped layered high- T_c cuprate parent compounds which have a non-frustrated square lattice arrangement of the Cu^{+2} ions (see [O23]). We thus anticipate that if $\text{SrCu}_2(\text{BO}_3)_2$ can be doped, the evolutions in the magnetic and electronic properties with doping might be qualitatively different from those known for the layered high- T_c cuprates, offering the potential for interesting new physics and physical behaviors. We found that sintered pellets of the compound do not react significantly with water and therefore that aqueous chemical and electrochemical doping techniques were potentially possible. We verified literature reports [L43-L45] that La_2CuO_4 can be hole-doped to become superconducting by reacting it with aqueous sodium hypochlorite (NaClO) solution at room temperature for several days. We then used this technique to attempt to chemically oxidize $\text{SrCu}_2(\text{BO}_3)_2$. We found that both water and aqueous NaClO have little effect on the magnetic susceptibility of sintered pellets of $\text{SrCu}_2(\text{BO}_3)_2$ as shown in Fig. 8. Powdered samples of $\text{SrCu}_2(\text{BO}_3)_2$, however, partially decomposed to CuO and a water soluble strontium borate complex when exposed to water for several hours. The powdered samples completely decomposed to CuO and SrCO_3 when exposed to the NaClO solution (the CO_2 in the SrCO_3 came from the air and CO_2 dissolved in the water). Magnetic susceptibility data on the powder




Fig. 7 (Left figure) The structure of $\text{SrCu}_2(\text{BO}_3)_2$ viewed along the c axis. The box shows a unit cell projected on the ab plane. The Sr atoms are displaced above and below the ab plane.

Fig. 8 (Right figure) (Color) Magnetic Susceptibility χ versus temperature T in an applied magnetic field of $H = 10$ kOe for pellets of as-made $\text{SrCu}_2(\text{BO}_3)_2$ (\diamond), $\text{SrCu}_2(\text{BO}_3)_2$ reacted with water (squares), $\text{SrCu}_2(\text{BO}_3)_2$ reacted with aqueous NaClO for 12 h (circles), and $\text{SrCu}_2(\text{BO}_3)_2$ reacted with NaClO for 72 h (triangles) [P44]. The “as-made” sample is one that received no special treatments after synthesis.

samples were both quantitatively and qualitatively different from the magnetic susceptibility of as-prepared $\text{SrCu}_2(\text{BO}_3)_2$ and consistent with the presence of a large concentration of CuO in the sample.

In addition to the aqueous chemical doping experiments discussed above, we tried to partially replace the Sr in $\text{SrCu}_2(\text{BO}_3)_2$ by La and fire the samples in air, and then planned to anneal the samples in an inert atmosphere to remove excess oxygen and thereby electron dope the material. The strategy was the same as that successfully used in the past to synthesize the electron-doped high T_c cuprate $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ [O23]. We found that LaBO_3 formed as a second phase in all samples containing La, the lattice parameters of the $\text{SrCu}_2(\text{BO}_3)_2$ were invariant with the amount of La doping, and that the magnetic susceptibility of those samples was qualitatively unchanged from the magnetic susceptibility of pure $\text{SrCu}_2(\text{BO}_3)_2$. These results all indicated that when synthesized in air, excess oxygen does not enter the lattice to form $\text{SrCu}_2(\text{BO}_3)_2\text{O}_x$, in contrast to the cuprate $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4+x}$ that is formed in air. Additional electron doping experiments are in progress to synthesize $\text{Sr}_{1-x}\text{La}_x\text{Cu}_2(\text{BO}_3)_2$ in sealed quartz tubes. We are also planning to synthesize $\text{Sr}_{1-x-y}\text{La}_x\text{Ba}_y\text{Cu}_2(\text{BO}_3)_2$ in which x and y will be adjusted to keep the average (La, Ba) cation size the same as for Sr in order to reduce lattice mismatch effects. Finally, we are attempting to hole-dope $\text{SrCu}_2(\text{BO}_3)_2$ by substituting Na^{+1} for Sr^{+2} .

MgAlB₁₄

The recent observation of superconductivity below a temperature of 39 K in the previously known compound MgB_2 [L46] indicated that the condensed matter physics community may have overlooked interesting physical properties in other known non-transition metal compounds. The boride compound MgAlB_{14} was discovered by Ames Lab scientists a few years ago to be one of the hardest materials known [L47]. The crystal structure of MgAlB_{14} consists of linear (zig-zag) chains of Al (Mg), but crystallographic studies in the literature report that the metal sites are not fully occupied; the occupation of the cation sites (0.8) is such that the cations together transfer four electrons to the boron sublattice [L48-L50]. In addition to MgAlB_{14} , the compound Mg_2B_{14} [L51] forms in the same structure as well as MAlB_{14} where $M = \text{Li, Y, Tb, Dy, Ho, Er, Tm, Yb, and Lu}$ [L49, L52-L54]. Only the Mg_2B_{14} and LiAlB_{14} compounds have fully occupied metal sites, again strongly indicating that the

compound formation is dependent on electronic effects. We completed a search for unusual magnetic properties in MgAlB_{14} in collaboration with B. A. Cook, J. L. Harringa, and A. M. Russell of Ames Laboratory [229]. Two types of samples were synthesized and studied: 1) powder and 2) chemically substituted and unsubstituted hot pressed pellets prepared from mechanically alloyed powders. We found that synthesis of single-phase powder samples of MgAlB_{14} was not possible, mostly due to the difficulty of incorporating the Al metal starting material [P48]. Magnetic susceptibility χ versus temperature T , magnetization M versus T , and isothermal M versus magnetic field H studies of our samples were carried out to search for superconductivity or ferromagnetism in this compound. The $\chi(T)$ measurements on a powder sample revealed temperature-independent diamagnetism with a small Curie-Weiss impurity concentration equivalent to about 1 mol% of spin-1/2 ions. In contrast, $M(T)$ and $M(H)$ data on the hot pressed samples showed evidence of ferromagnetic transitions above about 330 K. Scanning electron microscopy and Auger microprobe analysis of the hot pressed samples indicated that the samples contained significant concentrations of Fe impurities. We conclude that pure MgAlB_{14} is neither a superconductor nor a ferromagnet above 1.8 K and exhibits temperature-independent diamagnetism from 1.8 K up to room temperature. The ferromagnetism observed in the hot pressed samples is likely due to Fe impurities abraded from the stainless steel mills used to mix the starting materials prior to hot pressing the samples.

Outside References

- [L1]. “Possible High T_c Superconductivity in the Ba-La-Cu-O System”, J. G. Bednorz and K. A. Müller, *Z. Phys. B* **64**, 189-193 (1986).
- [L2]. “Bulk Superconductivity at 36 K in $\text{La}_{1.8}\text{Sr}_{0.2}\text{CuO}_4$ ”, R. J. Cava, R. B. van Dover, B. Batlogg, and E. A. Rietman, *Phys. Rev. Lett.* **58**, 408-410 (1987).
- [L3]. “Superconductivity in Alkaline Earth-Substituted $\text{La}_2\text{CuO}_{4-y}$ ”, J. Georg Bednorz, K. Alex Müller, and Masaaki Takashige, *Science* **236**, 73-75 (1987).
- [L4]. “Antiferromagnetic Spin Correlations in Insulating, Metallic, and Superconducting $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ”, R. J. Birgeneau, D. R. Gabbe, H. P. Jenssen, M. A. Kastner, P. J. Picone, T. R. Thurston, G. Shirane, Y. Endoh, M. Sato, K. Yamada, Y. Hidaka, M. Oda, Y. Enomoto, M. Suzuki, and T. Murakami, *Phys. Rev. B* **38**, 6614-6623 (1988).
- [L5]. “Evidence for Stripe Correlations of Spins and Holes in Copper Oxide Superconductors”, J. M. Tranquada, B. J. Sternlieb, J. D. Axe, Y. Nakamura, and S. Uchida, *Nature* **375**, 561-563 (1995).
- [L6]. D. B. Rogers, J. L. Gillson, and T. E. Gier, *Solid State Commun.* **5**, 263 (1967).
- [L7]. H. Kessler and M. J. Sienko, *J. Chem. Phys.* **55**, 5414 (1971).
- [L8]. “Transport Properties of Metallic LiV_2O_4 Single Crystals-Heavy Mass Fermi Liquid Behavior”, H. Takagi, C. Urano, S. Kondo, M. Nohara, Y. Ueda, T. Shiraki, and T. Okubo, *Mater. Sci. Eng.* **B63**, 147-150 (1999).
- [L9]. “From the Geometrically Frustrated Antiferromagnets ZnV_2O_4 and ZnCr_2O_4 to the Heavy-Mass Fermi Liquid LiV_2O_4 ”, Shinichiro Kondo, Chiharu Urano, Yusuke Kurihara, Minoru Nohara, and Hidenori Takagi, *J. Phys. Soc. Jpn.* **69**, Suppl. B, 139-143 (2000).
- [L10]. “ LiV_2O_4 Spinel as a Heavy-Mass Fermi Liquid: Anomalous Transport and the Role of Frustration”, C. Urano, M. Nohara, S. Kondo, F. Sakai, H. Takagi, T. Shiraki, and T. Okubo, *Phys. Rev. Lett.* **85**, 1052-1055 (2000).
- [L11]. “Dramatic Change of the Magnetic Response in LiV_2O_4 : Possible Heavy Fermion to Itinerant d -Metal Transition”, A. Krimmel, A. Loidl, M. Klemm, S. Horn, and H. Schober, *Phys. Rev. Lett.* **82**, 2919-2922 (1999).
- [L12]. “Comment on ‘Dramatic Change of the Magnetic Response in LiV_2O_4 : Possible Heavy Fermion to Itinerant d -Metal Transition’”, A. P. Murani, *Phys. Rev. Lett.* **85**, 3981 (2000).
- [L13]. “Krimmel et al. Reply”, A. Krimmel, A. Loidl, M. Klemm, S. Horn, and H. Schober, *Phys. Rev. Lett.* **85**, 3982 (2000).
- [L14]. “Spin Fluctuations in a Magnetically Frustrated Metal LiV_2O_4 ”, S.-H. Lee, Y. Qiu, C. Broholm, Y. Ueda, and J. J. Rush, *Phys. Rev. Lett.* **86**, 5554-5557 (2001).
- [L15]. “Heavy Fermions in LiV_2O_4 : Kondo Compensation versus Geometric Frustration”, H. Kaps, M. Brando, W. Trinkl, N. Büttgen, A. Loidl, W.-W. Scheidt, M. Klemm, and S. Horn, *J. Phys. Condens. Mat.* **13**, 8497-8508 (2001).
- [L16]. “Ground State Properties of LiV_2O_4 and $\text{Li}_{1-x}\text{Zn}_x(\text{V}_{1-y}\text{Ti}_y)\text{O}_4$ ”, M. Brando, N. Büttgen, V. Fritsch, J. Hemberger, H. Kaps, H.-A. Krug von Nidda, M. Nicklas, K. Pucher, W. Trinkl, A. Loidl, E. W. Scheidt, M. Klemm, and S. Horn, *Eur. Phys. J. B* **25**, 289-298 (2002).
- [L17]. “Magnetic Correlations in Frustrated LiV_2O_4 and ZnV_2O_4 ”, N. Büttgen, A. Krimmel, A. Loidl, M. Klemm, S. Horn, D. R. Noakes, E. Schreier, and G. M. Kalvius, *Physica B* **312-313**, 703-705 (2002).
- [L18]. “A New Compound, $\text{Lu}_2\text{V}_2\text{O}_7$ ”, K. Kitayama and T. Katsura, *Chem. Lett.*, 815-816 (1976).
- [L19]. “New Ferromagnetic Semiconductor Oxides with a Pyrochlore Structure”, G. Bazuev, D. Makarova, V. Oboldin, and G. Shveikin, *Dokl. Akad. Nauk SSSR* **230**, 869-871 (1976).

- [L20]. “Magnetic Properties of Rare Earth Element Hypovanadates with Pyrochlore Structure“, G. Bazuev, A. Samokhvalov, Y. Morozov, I. Matveenkov, V. Babushkin, T. Arbuzova, and G. Shveikin, *Fizika Tverdogo Tela* **19**, 3274-3278 (1977).
- [L21]. “Preparation and Crystal Chemistry of Some Rare Earth Vanadium (IV) Pyrochlores $(RE)_2V_2O_7$; RE = Lu, Yb, Tm, $Lu_{1-x}Y_x$, $Lu_{1-x}Sc_x$ “, J. Greedan, *Mater. Res. Bull.* **14**, 13-19 (1979).
- [L22]. “Magnetic Properties of the Pyrochlores $(RE)_2V_2O_7$; RE = Tm, Yb, Lu. New Ferri- and Ferromagnetic Semiconductors“, L. Soderholm and J. Greedan, *Rare Earths in Modern Science and Technology* **2**, 393-4 (1980).
- [L23]. “Low-Temperature Behavior of Two-Dimensional Quantum Antiferromagnets“, Sudip Chakravarty, Bertrand I. Halperin, and David R. Nelson, *Phys. Rev. Lett.* **60**, 1057-1060 (1988).
- [L24]. “Two-Dimensional Quantum Heisenberg Antiferromagnet at Low Temperatures“, Sudip Chakravarty, Bertrand I. Halperin, and David R. Nelson, *Phys. Rev. B* **39**, 2344-2371 (1989).
- [L25]. “Susceptibility of the Spin 1/2 Heisenberg Antiferromagnetic Chain“, Sebastian Eggert, Ian Affleck, and Minoru Takahashi, *Phys. Rev. Lett.* **73**, 332-335 (1994).
- [L26]. “Linear Magnetic Chains with Anisotropic Coupling“, Jill C. Bonner and Michael E. Fisher, *Phys. Rev.* **135**, A640-A658 (1964).
- [L27]. “Reduction of Ordered Moment and Néel Temperature of Quasi-One-Dimensional Antiferromagnets Sr_2CuO_3 and Ca_2CuO_3 “, K. M. Kojima, Y. Fudamoto, M. Larkin, G. M. Luke, J. Merrin, B. Nachumi, Y. J. Uemura, N. Motoyama, H. Eisaki, S. Uchida, K. Yamada, Y. Endoh, S. Hosoya, B. J. Sternlieb, and G. Shirane, *Phys. Rev. Lett.* **78**, 1787-1790 (1997).
- [L28]. “Crystal Structure of Strontium Hydroxide [Copper Hydroxide] $Sr_2(OH)_2[Cu(OH)_4]$ “, T. N. Nadezhina, E. A. Pobedinskaya, and N. V. Belov, *Dokl. Akad. Nauk SSSR* **250**, 1126-1130 (1980) [*Sov. Phys. Dokl.* **25**, 73-75 (1980)]; “Einkristalle und Struktur von Hydroxocupraten“, E. Dubler, P. Korber, and H. R. Oswald, *Naturwissenschaften* **59**, 467-468 (1972); “The Crystal Structures of Barium Hexahydroxocuprate(II) and Strontium Hexahydroxocuprate(II), $Ba_2[Cu(OH)_6]$ and $Sr_2[Cu(OH)_6]$ “, E. Dubler, P. Korber, and H. R. Oswald, *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.* **29**, 1929-1933 (1973).
- [L29]. “Evidence of Oxygen Intercalation and Mobility at Room Temperature in Oxides: An Electrochemical Quartz Microbalance Study of Intercalation in La_2CuO_4 “, N. Casañ-Pastor, C. Zinck, C. R. Michel, E. M. Tejada-Rosales, and G. Torres-Gómez, *Chem. Mater.* **13**, 2118-2126 (2001).
- [L30]. “Magnetic Excitations in the $S = 1/2$ Alternating Chain Compound $(VO)_2P_2O_7$ “, A. W. Garrett, S. E. Nagler, D. A. Tennant, B. C. Sales, and T. Barnes, *Phys. Rev. Lett.* **79**, 745-748 (1997).
- [L31]. T. M. Rice, S. Gopalan, and M. Sigrist, *Europhys. Lett.* **23**, 445 (1993).
- [L32]. “Spin Ladders with Spin Gaps: A Description of a Class of Cuprates“, Sudha Gopalan, T. M. Rice, and M. Sigrist, *Phys. Rev. B* **49**, 8901-8910 (1994).
- [L33]. “Observation of a Spin Gap in $SrCu_2O_3$ Comprising Spin-1/2 Quasi-1D Two-Leg Ladders“, M. Azuma, Z. Hiroi, M. Takano, K. Ishida, and Y. Kitaoka, *Phys. Rev. Lett.* **73**, 3463-3466 (1994).
- [L34]. “Crystal Structure and Spin Gap State of CaV_2O_5 “, M. Onoda and N. Nishiguchi, *J. Solid State Chem.* **127**, 359-362 (1996).
- [L35]. “Magnetic Interaction in Insulating Cuprates“, Yoshiaki Mizuno, Takami Tohyama, and Sadamichi Maekawa, *J. Low Temp. Phys.* **117**, 389-393 (1999).

- [L36]. "Spin Waves and Electronic Interactions in La_2CuO_4 ", R. Coldea, S. M. Hayden, G. Aeppli, T. G. Perring, C. D. Frost, T. E. Mason, S.-W. Cheong, and Z. Fisk, *Phys. Rev. Lett.* **86**, 5377-5380 (2001).
- [L37]. "Dependence of Lattice Parameters and T_c on the Hole Concentration Determined by Precise Measurements of the Oxygen Content of $\text{M}_x\text{La}_{2-x}\text{CuO}_{4-y}$ ($\text{M} = \text{Ba}, \text{Sr}, \text{Ca}$)", K. Oh-ishi and Y. Syono, *J. Solid State Chem.* **95**, 136-144 (1991).
- [L38]. "Exact Dimer Ground State and Quantized Magnetization Plateaus in the Two-Dimensional Spin System $\text{SrCu}_2(\text{BO}_3)_2$ ", H. Kageyama, K. Yoshimura, R. Stern, N. V. Mushnikov, K. Onizuka, M. Kato, K. Kosuge, C. P. Slichter, T. Goto, and Y. Ueda, *Phys. Rev. Lett.* **82**, 3168-3171 (1999).
- [L39]. "Exact Dimer Ground State of the Two Dimensional Heisenberg Spin System $\text{SrCu}_2(\text{BO}_3)_2$ ", S. Miyahara and K. Ueda, *Phys. Rev. Lett.* **82**, 3701-3704 (1999).
- [L40]. "Exact Ground State of a Quantum Mechanical Antiferromagnet", B. S. Shastry and B. Sutherland, *Physica B* **108**, 1069-1070 (1981).
- [L41]. "Synthesis, Structure, and Properties of the Orthoborate $\text{SrCu}_2(\text{BO}_3)_2$ ", R. W. Smith and D. A. Keszler, *J. Solid State Chem.* **93**, 430-435 (1991).
- [L42]. " $\text{SrCu}_2(\text{BO}_3)_2$: A Unique Mott Hubbard Insulator", B. S. Shastry and B. Kumar, *Prog. Theor. Phys. Suppl.* **145**, 1-16 (2002).
- [L43]. "Superconductivity at 45 K in $\text{La}_2\text{CuO}_{4+x}$ Oxidized by NaClO ", Q. Y. Tu, X. L. Chen, B. K. Ma, Z. X. Zhao, Y. C. Lan, and J. K. Liang, *Appl. Phys. A* **74**, 79-82 (2002).
- [L44]. "Structural and Superconducting Properties of $\text{La}_2\text{CuO}_{4+x}$ Chemically Oxidized by NaClO ", Q. Y. Tu, X. L. Chen, B. K. Ma, Z. X. Zhao, J. Q. Li, and J. K. Liang, *Physica C* **370**, 94-100 (2002).
- [L45]. "Study on the Structure and Superconductivity of $\text{La}_2\text{CuO}_{4+x}$ Chemically Oxidized by NaClO at 50 °C", Q. Y. Tu, X. L. Chen, B. K. Ma, Z. X. Zhao, Y. P. Xu, B. Q. Hu, and J. K. Liang, *Supercond. Sci. Technol.* **14**, 517-521 (2001).
- [L46]. "Superconductivity at 39 K in Magnesium Diboride", J. Nagamatsu, N. Nakagawa, T. Muranaka, Y. Zenitani, and J. Akimitsu, *Nature* **410**, 63-64 (2001).
- [L47]. "A New Class of Ultra-Hard Materials Based on AlMgB_{14} ", B. A. Cook, J. L. Haringa, T. L. Lewis, and A. M. Russell, *Scr. Mater.* **42**, 597-602 (2000).
- [L48]. "Refinement of the Structure of MgAlB_{14} ", I Higashi and T. Ito, *J. Less-Common Met.* **92**, 239-246 (1983).
- [L49]. "Boron-rich Crystals in Al-M-B ($\text{M} = \text{Li}, \text{Be}, \text{Mg}$) Systems Grown from High-Temperature Aluminum Solutions", I. Higashi, M. Kobayashi, S. Okada, K. Hamano, and T. Lundström, *J. Cryst. Growth* **128**, 1113-1119 (1993).
- [L50]. "Structure of MgAlB_{14} and a Brief Critique of Structural Relationships in Higher Borides", V. I. Matkovich and J. Economy, *Acta Crystallogr. Sect. B* **26**, 616-621 (1970).
- [L51]. "Crystal Structure of Magnesium Heptaboride Mg_2B_{14} ", A. Guette, M. Barret, R. Naslain, and P. Hagenmuller, *J. Less-Common Met.* **82**, 325-334 (1981).
- [L52]. "Crystal Structure, Electrical, and Magnetic Properties of the New Ternary Compounds LnAlB_{14} ", M. M. Korsukova, V. N. Gurin, Y. B. Kuzma, N. F. Chaban, S. I. Chykhrii, V. V. Moshchalkov, N. B. Brandt, A. A. Gippius, and K. Khyu Nyan, *Phys. Stat. Sol. A* **114**, 265-272 (1989).
- [L53]. "Crystal Structural Refinement of the New Compound TmAlB_{14} ", M. M. Korsukova, V. N. Gurin, Y. Yu, L.-E. Tergenius, and T. Lundström, *J. Alloys Compd.* **190**, 185-187 (1993).
- [L54]. "The Crystal Structure of Defective YAlB_{14} and ErAlB_{14} ", M. M. Korsukova, T. Lundström, L.-E. Tergenius, and V. N. Gurin, *J. Alloys Compd.* **187**, 39-48 (1992).

OTHER PUBLICATIONS (Not listed below in "Refereed Publications")

- O1. "Superconducting and Normal State Properties of Several Ternary Titanium Oxides," D. C. Johnston, Ph. D. Thesis, University of California at San Diego (March 1975), 207 pp.; available from University Microfilms, 300 N. Zeeb Road, Ann Arbor, MI 48106: Order #75-11,230; Dissertation Abstracts International 35 (11) (1975).
- O2. "Calorimetric Observation of a Phase Transition in the Superconducting State in $\text{Gd}_{1.2}\text{Mo}_6\text{Se}_8$," R. W. McCallum, D. C. Johnston, R. N. Shelton, and M. B. Maple, in Superconductivity in d- and f-Band Metals, Second Rochester Conference, D. H. Douglass, editor (Plenum, New York, 1976), pp. 625-634.
- O3. "Kondo Anomalies Due to Rare Earth Impurity Ions," M. B. Maple, L. E. DeLong, W. A. Fertig, D. C. Johnston, R. W. McCallum, and R. N. Shelton, in Valence Instabilities and Related Narrow-Band Phenomena, edited by R. D. Parks (Plenum, New York, 1977), pp. 17-29.
- O4. "Pressure Dependencies of the Superconducting and Magnetic Critical Temperatures of Ternary Rhodium Borides," R. N. Shelton and D. C. Johnston, in High-Pressure Physics, edited by C. W. Chu and J. A. Woollam (Plenum Press, New York, 1978), pp. 409-418.
- O5. "Neutron Scattering Study of the Magnetic Transition in the Re-entrant Superconductor $\text{Ho}_{0.6}\text{Er}_{0.4}\text{Rh}_4\text{B}_4$," H. A. Mook, W. C. Koehler, M. B. Maple, Z. Fisk, and D. C. Johnston, in Proc. Third Internat. Conf. on d- and f-band Metals, H. Suhl and M. B. Maple, editors, (Academic Press, New York, 1980), pp. 427-432.
- O6. "Crystal Field Effects in the Magnetic Susceptibility of Dilute Alloys of LaSn_3 Containing Nd Impurities," L. E. DeLong, M. Tovar, M. B. Maple, and D. C. Johnston, in Proc. Internat. Conf. on Crystalline Electric Field and Structural Effects in f-Electron Systems, Philadelphia, Pennsylvania, November 12-15, 1979, J. E. Crow, R. P. Guertin, T. W. Mihalison, ed. (Plenum, New York, 1980), pp. 581-588.
- O7. "Ultrasonic Attenuation Study of Induced Magnetic Order in ErRh_4B_4 ," S. Schneider, R. Chen, M. Levy, D. C. Johnston, and B. T. Matthias, in Ternary Superconductors, ed. by G. K. Shenoy, B. D. Dunlap, and F. Y. Fradin (Elsevier North Holland, New York, 1981), pp. 147-150.
- O8. "Precision Magnetization and Susceptibility Measurements on ErRh_4B_4 in the Ferromagnetic and Superconducting Phases," F. Behroozi, G. W. Crabtree, S. A. Campbell, M. Levy, D. Snider, D. C. Johnston, and B. T. Matthias, *ibid.*, pp. 155-158.
- O9. "Neutron Scattering Studies of the Magnetic Transitions in the Ternary Alloy System $\text{Er}_{1-x}\text{Ho}_x\text{Rh}_4\text{B}_4$," H. A. Mook, M. B. Maple, Z. Fisk, D. C. Johnston, and L. D. Woolf, *ibid.*, pp. 179-180.
- O10. "Effect of Pressure on the Superconducting and Magnetic Critical Temperatures of bct Ternary Ruthenium Borides," R. N. Shelton, C. U. Segre, and D. C. Johnston, *ibid.*, pp. 205-208.

OTHER PUBLICATIONS, CONT'D (Not listed in "Refereed Publications")

- O11. "Systematics of Superconductivity in Ternary Compounds," D. C. Johnston and H. F. Braun, Chap. 2 in *Superconductivity in Ternary Compounds*, Vol. 2, ed. by Ø. Fischer and M. B. Maple (Springer-Verlag, Heidelberg, 1982).
- O12. "Variation in the Structural, Magnetic and Superconducting Properties of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ with Oxygen Content," D. C. Johnston, A. J. Jacobson, J. M. Newsam, J. T. Lewandowski, and D. P. Goshorn, Ch. 14 in *Chemistry of High-Temperature Superconductors*, ACS Symposium Series Vol. 351, ed. by D. L. Nelson, M. S. Whittingham and T. F. George (Am. Chem. Soc., Washington, DC, 1987), pp. 136-151.
- O13. "Preparation of $\text{VO}(\text{HPO}_4) \cdot 0.5 \text{H}_2\text{O}$ and its Transformation to the Maleic Anhydride Catalyst $(\text{VO})_2\text{P}_2\text{O}_7$," J. W. Johnson, D. C. Johnston, and A. J. Jacobson, in *Preparation of Catalysts IV*, ed. by B. Delmon, P. Grange, P. A. Jacobs, and G. Poncelet (Elsevier, Amsterdam, 1987), pp. 181-190.
- O14. "Synthesis, Structural Chemistry and Properties of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$," A. J. Jacobson, J. M. Newsam, D. C. Johnston, J. P. Stokes, S. Bhattacharya, J. T. Lewandowski, D. P. Goshorn, M. J. Higgins and M. S. Alvarez, in *Chemistry of Oxide Superconductors*, ed. C. N. R. Rao (IUPAC/Blackwell, 1988), pp. 43-61.
- O15. "Magnetic Susceptibility Anisotropy in the High T_c Cuprates", D. C. Johnston, T. Matsumoto, Y. Yamaguchi, Y. Hidaka, and T. Murakami, in *Electronic Properties and Mechanisms of High T_c Superconductors*, edited by T. Oguchi, K. Kadowaki and T. Sasaki (Elsevier, Amsterdam, 1992), pp. 301-306.
- O16. "The Phase Diagrams and Doped-Hole Segregation in $\text{La}_2\text{CuO}_{4+y}$ and $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4+y}$ ($x \leq 0.15$, $y \leq 0.12$)", D. C. Johnston, F. Borsa, P. C. Canfield, J. H. Cho, F. C. Chou, L. L. Miller, D. R. Torgeson, D. Vaknin, J. Zarestky, J. Ziolo, J. D. Jorgensen, P. G. Radaelli, A. J. Schultz, J. L. Wagner, S-W. Cheong, W. R. Bayless, J. E. Schirber and Z. Fisk, in *Phase Separation in Cuprate Superconductors*, edited by E. Sigmund and K. A. Müller (Springer-Verlag, Heidelberg, 1994), pp. 82-100.
- O17. "Superconductivity, Magnetism and Metal-Insulator Transitions in Some Ternary and Pseudoternary 3d-, 4d- and 5d-Metal Oxides," D. C. Johnston, T. Ami, F. Borsa, M. K. Crawford, J. A. Fernandez-Baca, K. H. Kim, R. L. Harlow, A. V. Mahajan, L. L. Miller, M. A. Subramanian, D. R. Torgeson, and Z. R. Wang, in *Spectroscopy of Mott Insulators and Correlated Metals*, edited by A. Fujimori and Y. Tokura, Springer Series in Solid-State Sciences, Vol. 119 (Springer-Verlag, Berlin and Heidelberg, 1995), pp. 241-254.
- O18. "Anisotropic Magnetization of $\text{HoNi}_2\text{B}_2\text{C}$ ", P. C. Canfield, B. K. Cho, D. C. Johnston, and M. F. Hundley, *Mater. Res. Soc. Symp. Proc.* 376, 565-570 (1995).
- O19. "Magnetic Structures in $\text{RNi}_2\text{B}_2\text{C}$ (R=Ho, Er) Superconductors", C. Stassis, A. I. Goldman, P. Dervenis, J. Zarestky, P. C. Canfield, B. K. Cho, D. C. Johnston, and B. Sternlieb, *Mater. Res. Soc. Symp. Proc.* 376, 559-564 (1995).

OTHER PUBLICATIONS, CONT'D (Not listed in "Refereed Publications")

- O20. "Overview of Recent Magnetic Studies of High T_c Cuprate Parent Compounds and Related Materials", D. C. Johnston, T. Ami, F. Borsa, P. C. Canfield, P. Carretta, B. K. Cho, J. H. Cho, F. C. Chou, M. Corti, M. K. Crawford, P. Dervenagas, R. W. Erwin, J. A. Fernandez-Baca, A. I. Goldman, R. J. Gooding, Q. Huang, M. F. Hundley, R. L. Harlow, B. N. Harmon, A. Lascialfari, L. L. Miller, J. E. Ostenson, N. M. Salem, C. Stassis, B. Sternlieb, B. J. Suh, D. R. Torgeson, D. Vaknin, K. J. E. Vos, X.-L. Wang, Z. R. Wang, M. Xu, and J. Zarestky, Proceedings of the First Polish-U.S. Conference on High Temperature Superconductivity, Wroclaw and Duszynki-Zdroj, Poland, 11-15 September 1995, in Lecture Notes in Physics, Vol. 475, "Recent Developments in High Temperature Superconductivity" (Springer, Berlin and Heidelberg, 1996), pp. 61-82.
- O21. "Magnetic and Structural Properties and Phase Diagrams of $Sr_2CuO_2C_{12}$ and Lightly-Doped $La_{2-x}Sr_xCuO_{4+y}$ ", D. C. Johnston, F. Borsa, J. H. Cho, L. L. Miller, B. J. Suh, D. R. Torgeson, P. Carretta, M. Corti, A. Lascialfari, R. J. Gooding, N. M. Salem, K. J. E. Vos, and F. C. Chou, Proc. 10th Anniversary High Temperature Superconductivity Workshop on Physics, Materials and Applications, Houston, TX, 12-16 March 1996, edited by B. Batlogg, C. W. Chu, W. K. Chu, D. U. Gubser and K. A. Müller (World Scientific, Singapore, 1996), pp. 373-376.
- O22. "Magnetic Properties of Single-Layer Cuprates", D. C. Johnston, F. Borsa, P. Carretta, J. H. Cho, F. C. Chou, M. Corti, R. J. Gooding, E. Lai, A. Lascialfari, L.L. Miller, N. M. Salem, B. J. Suh, D. R. Torgeson, D. Vaknin, K.J.E. Vos, and J. L. Zarestky, in High- T_c Superconductivity 1996: Ten Years After the Discovery, edited by E. Kaldis, E. Liarokapis, and K. A. Müller (Kluwer, Amsterdam, 1997), pp. 311-348.
- O23. "Normal-State Magnetic Properties of Single-Layer Cuprate High-Temperature Superconductors and Related Materials", D. C. Johnston, in Handbook of Magnetic Materials, Vol. 10, edited by K.H.J.Buschow, Ch. 1 (Elsevier, Amsterdam 1997), pp. 1-237; a review with 120 figures and 733 references.
- O24. "The Mixed State Properties of Superconducting YNi_2B_2C ", M.-I. Mun, B. K. Cho, S.-I. Lee, P. C. Canfield, and D. C. Johnston, Physica B 259-261, 599-600 (1999).

REFEREED PUBLICATIONS IN PRINT

A. JOURNAL ARTICLES

1. "A Quantitative Transmission Line Experiment," D. C. Johnston and B. G. Silbernagel, *Am. J. Phys.* **37**, 1207-1211 (1969).
2. "Superconductivity in the Pseudo-Binary A-15 Compounds $\text{Mo}_3\text{Os}_{1-x}\text{Ru}_x$ and $\text{Mo}_3\text{Ir}_{1-x}\text{Ru}_x$," D. C. Johnston, *Solid State Commun.* **11**, 1751-1754 (1972).
3. " $\text{Zr}_{3+x}\text{S}_4$: A New Superconducting Binary Sulfide," D. C. Johnston and A. Moodenbaugh, *Phys. Lett.* **41A**, 447-448 (1972).
4. "Stabilization of the Cubic A-15 Phase of Nb_3Sn ," R. Viswanathan and D. C. Johnston, *Mater. Res. Bull.* **8**, 589-592 (1973).
5. "High Temperature Superconductivity in the Li-Ti-O Ternary System," D. C. Johnston, H. Prakash, W. H. Zachariasen, and R. Viswanathan, *Mat. Res. Bull.* **8**, 777-784 (1973).
6. "Magnetic and Superconducting Properties of Niobium Oxides," H. R. Khan, Ch. J. Raub, W. E. Gardner, W. A. Fertig, D. C. Johnston, and M. B. Maple, *Mater. Res. Bull.* **9**, 1129-1135 (1974).
7. "Superconductivity in Two NaCl Structure Compounds: $-\text{ZrP}$ and ScS_{1+x} ," A. R. Moodenbaugh, D. C. Johnston, and R. Viswanathan, *Mater. Res. Bull.* **9**, 1671-1675 (1974).
8. "Pressure Dependence of the Superconducting Transition Temperature for Ternary Molybdenum Sulfides," R. N. Shelton, A. C. Lawson and D. C. Johnston, *Mater. Res. Bull.* **10**, 297-302 (1975).
9. "Low Temperature Heat Capacity of TTF(TCNQ)," R. Viswanathan and D. C. Johnston, *J. Phys. Chem. Solids* **36**, 1093-1096 (1975).
10. "Isotope Effect in the Resistivity of Scandium Hydride," Z. Fisk and D. C. Johnston, *Phys. Letters* **53A**, 39-40 (1975).
11. "Thermodynamic Study of Structural and Superconducting Transitions in a V_3Si Single Crystal," R. Viswanathan and D. C. Johnston, *Phys. Rev. B* **13**, 2877-2879 (1976).
12. "Superconducting and Normal State Properties of $\text{Li}_{1+x}\text{Ti}_{2-x}\text{O}_4$ Spinel Compounds. I. Preparation, Crystallography, Superconducting Properties, Electrical Resistivity, Dielectric Behavior, Magnetic Susceptibility," D. C. Johnston, *J. Low Temp. Phys.* **25**, 145-175 (1976).
13. "Superconducting and Normal State Properties of $\text{Li}_{1+x}\text{Ti}_{2-x}\text{O}_4$ Spinel Compounds. II. Low Temperature Heat Capacity," R. W. McCallum, D. C. Johnston, C. A. Luengo, and M. B. Maple, *J. Low Temp. Phys.* **25**, 177-193 (1976).
13. "Low Temperature Heat Capacity of Small Nb_3Sn Polycrystals by ac Calorimetry," R. Viswanathan and D. C. Johnston, *J. Low Temp. Phys.* **25**, 1-8 (1976).

REFEREED PUBLICATIONS IN PRINT (cont'd)

15. "Electrical Resistivity and Magnetic Susceptibility of ThOs_2 ," A. C. Lawson, D. C. Johnston, and B. T. Matthias, *Phys. Letters* **56A**, 209-210 (1976).
16. "Specific Heat of a New Metastable Phase of Scandium-Chromium," R. W. McCallum, D. C. Johnston, M. B. Maple, and B. T. Matthias, *Mat. Res. Bull.* **11**, 781-786 (1976); **11**, 1354 (1976).
17. "Measurement of the Pressure Dependence of T_c for Superconducting Spinel Compounds," R. N. Shelton, D. C. Johnston, and H. Adrian, *Solid State Commun.* **20**, 1077-1080 (1976).
18. "Superconductivity of Neutron Irradiated Mo_3Os ," A. R. Sweedler, S. Moehlecke, R. H. Jones, R. Viswanathan and D. C. Johnston, *Solid State Commun.* **21**, 1007-1009 (1977); **23**, vii (1977).
19. "Superconducting and Normal State Properties of $\text{Ag}_{1-x}\text{Sn}_{1+x}\text{Se}_{2-y}$," D. C. Johnston and H. Adrian, *J. Phys. Chem. Solids* **38**, 355-365 (1977).
20. "Magnetic Properties of $\text{RE}_x\text{Mo}_6\text{Se}_8$ Compounds Between 0.7 and 295 K," D. C. Johnston and R. N. Shelton, *J. Low Temp. Phys.* **26**, 561-572 (1977).
20. "Pressure Dependence of T_c for the Shear-Stress-Induced Phase of V_3Si ," R. N. Shelton, D. C. Johnston and R. Viswanathan, *Mater. Res. Bull.* **12**, 133-136 (1977).
22. "Aspherical Coulomb Scattering of Conduction Electrons in PrB_6 ," Z. Fisk and D. C. Johnston, *Solid State Commun.* **22**, 359-362 (1977).
23. "Phase Transition in the Superconducting State in Gd-Rich Pseudoternary Compounds $\text{La}_{1-x}\text{Gd}_x\text{Mo}_6\text{Se}_8$," R. W. McCallum, D. C. Johnston, R. N. Shelton and M. B. Maple, *Solid State Commun.* **24**, 391-395 (1977).
24. "Superconductivity, Lattice Transformations, and Electronic Instabilities in $\text{Cu}_x\text{Mo}_3\text{S}_4$," D. C. Johnston, R. N. Shelton, and J. J. Bugaj, *Solid State Commun.* **21**, 949-953 (1977).
25. "Destruction of Superconductivity at the Onset of Long-Range Magnetic Order in the Compound ErRh_4B_4 ," W. A. Fertig, D. C. Johnston, L. E. DeLong, R. W. McCallum, M. B. Maple, and B. T. Matthias, *Phys. Rev. Letters* **38**, 987-990 (1977).
26. "Coexistence of Superconductivity and Long-Range Antiferromagnetic Order in the Compound $\text{Er}_x\text{Mo}_6\text{Se}_8$," R. W. McCallum, D. C. Johnston, R. N. Shelton, W. A. Fertig, and M. B. Maple, *Solid State Commun.* **24**, 501-505 (1977).
27. "Superconductivity in a New Ternary Structure Class of Boride Compounds," D. C. Johnston, *Solid State Commun.* **24**, 699-702 (1977).
28. "Low Temperature Specific Heat of the Magnetic Superconductor $\text{Gd}_{1.0}\text{Mo}_6\text{Se}_8$," L. J. Azevedo, W. G. Clark, C. Murayama, R. W. McCallum, D. C. Johnston, M. B. Maple, and R. N. Shelton, *J. Phys., Colloq. C6* **39**, 365-6 (1978).

REFEREED PUBLICATIONS IN PRINT (cont'd)

29. "Superconducting and Magnetic Properties of ErRh_4B_4 ," H. R. Ott, W. A. Fertig, D. C. Johnston, M. B. Maple, and B. T. Matthias, *J. Phys., Colloq. C6* **39**, 375-6 (1978).
30. "Re-entrant Superconductivity and Magnetic Ordering in the Pseudoternary System $(\text{Er}_{1-x}\text{Ho}_x)\text{Rh}_4\text{B}_4$," D. C. Johnston, W. A. Fertig, M. B. Maple and B. T. Matthias, *Solid State Commun.* **26**, 141-144 (1978).
31. "Electron Spin Resonance on $\text{Gd}_x\text{Mo}_6\text{Se}_8$ ($x = 1.0$ and 1.2)," S. Oseroff, R. Calvo, D. C. Johnston, M. B. Maple, R. W. McCallum and R. N. Shelton, *Solid State Commun.* **27**, 201-204 (1978).
32. "Superconducting and Magnetic Properties of ErRh_4B_4 ," H. R. Ott, W. A. Fertig, D. C. Johnston, M. B. Maple, and B. T. Matthias, *J. Low Temp. Phys.* **33**, 159-174 (1978).
33. "Superconductivity of Transition Metal Sulfides, Selenides and Phosphides with the NaCl Structure," A. R. Moodenbaugh, D. C. Johnston, R. Viswanathan, R. N. Shelton, L. E. DeLong, and W. A. Fertig, *J. Low Temp. Phys.* **33**, 175-203 (1978).
34. "Interaction Between Superconductivity and Magnetism in the Pseudoternary System $(\text{Lu}_{1-x}\text{Ho}_x)\text{Rh}_4\text{B}_4$," M. B. Maple, H. C. Hamaker, D. C. Johnston, H. B. MacKay, and L. D. Woolf, *J. Less Common Metals* **62**, 251-263 (1978).
35. "Superconductivity of Single Crystal NbGe_2 ," J. P. Remeika, A. S. Cooper, Z. Fisk, and D. C. Johnston, *J. Less Common Metals* **62**, 211-213 (1978).
36. "Anomalous Pressure Dependences of the Superconducting Transition Temperature of Dilute Alloys of LaSn_3 Containing Light Rare Earth Impurities," L. E. DeLong, M. B. Maple, R. W. McCallum, L. D. Woolf, R. N. Shelton, and D. C. Johnston, *J. Low Temp. Phys.* **34**, 445-485 (1979).
37. "Specific Heat Anomalies at the Lower Critical Temperature in Re-entrant Ferromagnetic Superconductors," H. B. MacKay, L. D. Woolf, M. B. Maple, and D. C. Johnston, *Phys. Rev. Lett.* **42**, 918-921 (1979); errata: 43, 89 (1979).
38. "Superconducting and Normal State Properties of ErRh_4B_4 and LuRh_4B_4 ," L. D. Woolf, D. C. Johnston, H. B. MacKay, R. W. McCallum, and M. B. Maple, *J. Low Temp. Phys.* **35**, 651-669 (1979).
39. "Magnetic, Transport, and Thermal Properties of Ferromagnetic EuB_6 ," Z. Fisk, D. C. Johnston, B. Cornut, S. von Molnar, S. Oseroff, and R. Calvo, *J. Appl. Phys.* **50**, 1911-1913 (1979).
40. "Superconductivity and Crystal Structure of Ternary $\text{Cu}_x\text{Mo}_3\text{S}_4$ Compounds with $0 \leq x \leq 1$ Prepared by Anodic Oxidation of $\text{Cu}_{1.0}\text{Mo}_3\text{S}_4$," M. Tovar, L. E. DeLong, D. C. Johnston, and M. B. Maple, *Solid State Commun.* **30**, 551-557 (1979).
41. "Magnetic and Electronic Properties of EuB_6 ," S. Oseroff, R. Calvo, J. Stankiewicz, Z. Fisk, and D. C. Johnston, *Phys. Stat. Sol. B* **94**, K133- K135 (1979).

REFEREED PUBLICATIONS IN PRINT (cont'd)

42. "ScRu₄B₄: A New Superconducting Member of the MRu₄B₄ Ternary System," H. C. Ku, D. C. Johnston, B. T. Matthias, H. Barz, G. Burri, and L. Rinderer, *Mater. Res. Bull.* **14**, 1591-1599 (1979).
43. "Magnetic Susceptibility of Nb₃Sb," G. W. Webb, Z. Fisk and D. C. Johnston, *Phys. Lett. A* **73**, 350-352 (1979).
44. "Superconducting and Normal State Properties of Dilute Alloys of LaSn₃ Containing Nd Impurities," L. E. DeLong, M. Tovar, L. D. Woolf, M. B. Maple, D. C. Johnston, and J. Keller, *J. Low Temp. Phys.* **38**, 119-147 (1980).
45. "¹¹B NMR Studies of the Electronic and Magnetic Properties of LuRh₄B₄ and GdRh₄B₄," D. C. Johnston and B. G. Silbernagel, *Phys. Rev. B* **21**, 4996-5003 (1980).
46. "Intercalation of Organometallic Compounds into Layered Transition Metal Oxyhalides," T. R. Halbert, D. C. Johnston, L. E. McCandlish, A. H. Thompson, J. C. Scanlon, and J. A. Dumesic, *Physica* **99B**, 128-132 (1980).
47. "Pressure Enhanced Superconductive and Magnetic Interactions in the System (Er_{1-x}Ho_x)Rh₄B₄," R. N. Shelton, C. U. Segre, and D. C. Johnston, *Solid State Commun.* **33**, 843-846 (1980).
48. "Low Temperature Magnetic Properties of HoRh₄B₄," H. R. Ott, L. D. Woolf, M. B. Maple, and D. C. Johnston, *J. Low Temp. Phys.* **39**, 383-396 (1980).
49. "Magnetic Field Dependence of the Neutron Scattering from ErRh₄B₄," H. A. Mook, M. B. Maple, Z. Fisk, and D. C. Johnston, *Solid State Commun.* **36**, 287-289 (1980).
50. "Superconducting and Magnetic Properties of New Ternary Borides with the CeCo₃B₂-type Structure," H. C. Ku, G. P. Meisner, F. Acker, and D. C. Johnston, *Solid State Commun.* **35**, 91-96 (1980).
51. "The Electronic Structure of Fe²⁺ in Reaction Centers from R. Sphaeroides. I. Static Magnetization Measurements," W. F. Butler, D. C. Johnston, H. B. Shore, D. R. Fredkin, M. Y. Okamura, and G. Feher, *Biophysical Journal* **32**, 967-992 (1980).
52. "Temperature Dependence of the Ultrasonic Attenuation Coefficient for ErRh₄B₄ between 1.5 K and 20 K," S. C. Schneider, M. Levy, D. C. Johnston, and B. T. Matthias, *Phys. Lett.* **80A**, 72-74 (1980).
53. "Ferromagnetism in the RERh₄B₄ Compounds," H. B. MacKay, L. D. Woolf, M. B. Maple and D. C. Johnston, *J. Low Temp. Phys.* **41**, 639-651 (1980).
54. "Oscillatory Magnetic Fluctuations Near the Superconductor-to-Ferromagnet Transition in ErRh₄B₄," D. E. Moncton, D. B. McWhan, P. H. Schmidt, G. Shirane, W. Thomlinson, M. B. Maple, H. B. MacKay, L. D. Woolf, Z. Fisk, and D. C. Johnston, *Phys. Rev. Lett.* **45**, 2060-2063 (1980).

REFEREED PUBLICATIONS IN PRINT (cont'd)

55. "Amorphous MoS₃ and WS₃," K. S. Liang, S. P. Cramer, D. C. Johnston, C. H. Chang, A. J. Jacobson, J. P. deNeufville, and R. R. Chianelli, *J. Non-Cryst. Solids* **42**, 345-356 (1980).
56. "Thermodynamic Critical Field of Superconducting ErRh₄B₄ as a Function of Temperature," F. Behroozi, M. Levy, D. C. Johnston, and B. T. Matthias, *Solid State Commun.* **38**, 515-519 (1981).
57. "Experimental Determination of the Magnetic Landau Parameters in ErRh₄B₄," F. Behroozi, G. W. Crabtree, S. A. Campbell, M. Levy, D. R. Snider, D. C. Johnston, and B. T. Matthias, *Solid State Commun.* **39**, 1041-1044 (1981).
58. "Nonmagnetic Contributions to the Variation of T_c with RE in RE(Rh,Ru)₄B₄Compounds," D. C. Johnston, *Physica* **108B**, 755-756 (1981).
59. "Low Temperature Thermal Conductivity of LuRh₄B₄ and ErRh₄B₄," W. Odoni, G. Keller, H. R. Ott, H. C. Hamaker, D. C. Johnston, and M. B. Maple, *Physica* **108B**, 1227-1228 (1981).
60. "Ultrasonic Attenuation Study of H_{c1}(T) for ErRh₄B₄," S. C. Schneider, M. Levy, M. Tachiki, and D. C. Johnston, *Physica* **108B**, 807-808 (1981).
61. "Ultrasonic Attenuation Determination of H_{c1} and H_{c2} for ErRh₄B₄ at 1.5K," S. C. Schneider, M. Levy, R. Chen, M. Tachiki, D. C. Johnston, and B. T. Matthias, *Solid State Commun.* **40**, 61-64 (1981).
62. "Hydrostatic Pressure Effects on the Superconducting Properties of Ag_{1-x}Sn_{1+x}Se₂," R. N. Shelton, C. U. Segre, and D. C. Johnston, *Solid State Commun.* **39**, 797-800 (1981).
63. "Multiple Phase Transitions in the Rare Earth Tetraborides at Low Temperature," Z. Fisk, M. B. Maple, D. C. Johnston, and L. D. Woolf, *Solid State Commun.* **39**, 1189-1192 (1981).
64. "Neutron Scattering Study of the Magnetic Transition in (Ho_{1-x}Er_x)Rh₄B₄ Alloys," H. A. Mook, W. C. Koehler, M. B. Maple, Z. Fisk, D. C. Johnston, and L. D. Woolf, *Phys. Rev. B* **25**, 372-380 (1982).
65. "HoRh₄B₄: A Model Mean-Field Ferromagnet," H. R. Ott, G. Keller, W. Odoni, L. D. Woolf, M. B. Maple, D. C. Johnston, and H. A. Mook, *Phys. Rev. B* **25**, 477-480 (1982).
66. "Ambient Temperature Phase Relations in the System Na_{1/3}(H₂O)_yTaS₂ (0 ≤ y ≤ 2)," D. C. Johnston, *Mat. Res. Bull.* **17**, 13-23 (1982).
67. "Orthorhombic LuRh₄B₄: A New Polytype of MT₄B₄ Stoichiometry," K. Yvon and D. C. Johnston, *Acta Cryst. B* **38**, 247-250 (1982).
68. "Interpretation of Specific Heat and Spontaneous Magnetization Anomalies at the Re-entrant Superconducting - Ferromagnetic Transition in (Ho_{0.6}Er_{0.4})Rh₄B₄," L. D. Woolf, D. C. Johnston, H. A. Mook, W. C. Koehler, M. B. Maple, and Z. Fisk, *Physica* **109 & 110B**, 2045-2047 (1982).

REFEREED PUBLICATIONS IN PRINT (cont'd)

69. "Observation of New Cointercalation Compounds in the System $\text{Na}_{1/3}(\text{H}_2\text{O})_y\text{TaS}_2$ ($0 \leq y \leq 2$) from Powder X-Ray Diffraction Measurements," D. C. Johnston, *J. Less-Common Metals* **84**, 327-347 (1982).
70. "Superconductivity and Magnetism in the $\text{Ho}_{1-x}\text{Er}_x\text{Rh}_4\text{B}_4$ Alloy System," H. A. Mook, W. C. Koehler, S. K. Sinha, G. W. Crabtree, D. G. Hinks, M. B. Maple, Z. Fisk, D. C. Johnston, L. D. Woolf, and H. C. Hamaker, *J. Appl. Phys.* **53**, 2614-2618 (1982).
71. "The Role of Boron in the Superconductivity of Ternary RT_4B_4 Compounds," D. C. Johnston, *Solid State Commun.* **42**, 453-456 (1982).
72. "Observation of the Antiferromagnetic Transition in the Linear Chain Compound KFeS_2 by Magnetic Susceptibility and Heat Capacity Measurements," D. C. Johnston, S. C. Mraw, and A. J. Jacobson, *Solid State Commun.* **44**, 255-258 (1982).
73. "Anisotropy of the Normal State Properties of the Superconducting $\text{Co}_{1-x}\text{Ni}_x\text{Zr}_2$ System," Z. Henkie, W. A. Fertig, Z. Fisk, D. C. Johnston, and M. B. Maple, *J. Low Temp. Phys.* **48**, 389-403 (1982).
74. "Magnetic Susceptibility and Band Structure of Intercalation Compounds of 2H-TaS_2 ," D. C. Johnston, *Solid State Commun.* **43**, 533-536 (1982).
75. "Characterization of the CDW Transition in Orthorhombic TaS_3 by Thermoelectric Power and Magnetic Susceptibility Measurements," D. C. Johnston, J. P. Stokes, P. L. Hsieh, and G. Grüner, *J. Physique* **44**, C3-1749 to C3-1752 (1983).
76. "Superconductivity and Magnetism in Ternary Rare-Earth Compounds," H. A. Mook, O. A. Pringle, S. Kawarazaki, S. K. Sinha, G. W. Crabtree, D. G. Hinks, M. B. Maple, Z. Fisk, D. C. Johnston, L. D. Woolf, and H. C. Hamaker, *Physica B+C* **120**, 197-201 (1983).
77. "X-ray Diffraction Study of $\text{Na}_{1/3}(\text{H}_2\text{O})_{1.5}\text{TaS}_2$: Observation of a Hendricks-Teller Disordered Layer Lattice," D. C. Johnston and S. P. Frysinger, *Phys. Rev. B* **30**, 980-984 (1984).
78. "Magnetic Susceptibility and ESR Measurements as Probes of Different Magnetic Species in MoS_3 ," D. C. Johnston, A. J. Jacobson, B. G. Silbernagel, S. P. Frysinger, S. M. Rich, and L. A. Gebhard, *J. Catal.* **52**, 2053-2056 (1984).
79. "Thermodynamics of Charge Density Waves in Quasi One-Dimensional Conductors," D. C. Johnston, *Phys. Rev. Lett.* **52**, 2053-2056 (1984).
80. "Preparation and Characterization of $\text{VOHPO}_4 \cdot 0.5\text{H}_2\text{O}$ and its Topotactic Transformation to $(\text{VO})_2\text{P}_2\text{O}_7$," J. W. Johnson, D. C. Johnston, A. J. Jacobson, and J. F. Brody, *J. Am. Chem. Soc.* **106**, 8123-8128 (1984).
81. "New Superconducting Ternary Transition Metal Compounds with the E9_3 -Type Structure," H. C. Ku and D. C. Johnston, *Chinese J. Phys.* **22**, 59-64 (1984).

REFEREED PUBLICATIONS IN PRINT (cont'd)

82. "Superconductivity and Magnetism of $M_x(H_2O)_yTaS_2$ Layered Cointercalation Compounds," D. C. Johnston and B. W. Keelan, *Solid State Commun.* **52**, 631-634 (1984).
83. "Influence of Charge Density Wave Fluctuations on the Magnetic Susceptibility of the Quasi One-Dimensional Conductor $(TaSe_4)_2I$," D. C. Johnston, M. Maki, and G. Grüner, *Solid State Commun.* **53**, 5-8 (1985).
84. "Synthesis and Characterization of the Cobalt Bis-tetra(thiomolybdate) Trianion, $Co(MoS_4)_3^{2-}$," W. H. Pan, D. C. Johnston, S. T. McKenna, R. R. Chianelli, T. R. Halbert, L. L. Hutchings, and E. I. Stiefel, *Inorg. Chem. Acta* **97**, L17-L20 (1985).
85. "Charge Density Wave Fluctuations in One Dimensional Metals," D. C. Johnston, *Solid State Commun.* **56**, 439-442 (1985).
86. "Formation of Ag_2S Under Ag Paint Electrodes in Contact with TaS_3 ," D. C. Johnston, J. P. Stokes, and R. A. Klemm, *Phys. Rev. B* **32**, 1330-1332 (1985).
87. "Spin Pairing and Variable Magnetic Defect Densities in the Maleic Anhydride Catalyst $(VO)_2P_2O_7$," D. C. Johnston and J. W. Johnson, *J. Chem. Soc., Chem. Commun.* **(23)**, 1720-1722 (1985).
88. "Relationships Between Magnetic Susceptibility and Single Particle Transport in Quasi-One-Dimensional Conductors," D. C. Johnston, J. P. Stokes, and R. A. Klemm, *J. Magn. Magn. Mater.* **54-57**, 1317-1318 (1986).
89. "Magnetic Field Induced Phases of $(TMTSF)_2ClO_4$," P. M. Chaikin, J. S. Brooks, R. V. Chamberlin, L. Y. Chiang, D. P. Goshorn, D. C. Johnston, M. J. Naughton, and X. Yan, *Physica* **143B**, 383-387 (1986).
90. "Magnetic Susceptibility of $(VO)_2P_2O_7$: A One-Dimensional Spin 1/2 Heisenberg Antiferromagnet with a Ladder Spin Configuration and a Singlet Ground State," D. C. Johnston, J. W. Johnson, D. P. Goshorn, and A. J. Jacobson, *Phys. Rev. B* **35**, 219-222 (1987).
91. "Synthesis and Study of Multiple Cationic Salts of Hexamethoxytriphenylene (HMTP)," L. Y. Chiang, D. C. Johnston, J. P. Stokes, and A. N. Bloch, *Synthetic Metals* **19**, 697-702 (1987).
92. "Antiferromagnetism in La_2CuO_{4-y} ," D. Vaknin, S. K. Sinha, D. E. Moncton, D. C. Johnston, J. M. Newsam, C. R. Safinya, and H. E. King, Jr., *Phys. Rev. Lett.* **58**, 2802-2805 (1987).
92. "Influence of Oxygen Defects on the Physical Properties of La_2CuO_{4-y} ," D. C. Johnston, J. P. Stokes, D. P. Goshorn, and J. T. Lewandowski, *Phys. Rev. B (Rapid Commun.)* **36**, 4007-4010 (1987).
94. "Confirmation of Antiferromagnetism in La_2CuO_{4-y} with Polarized Neutrons," S. Mitsuda, G. Shirane, S. K. Sinha, D. C. Johnston, M. S. Alvarez, D. Vaknin, and D. E. Moncton, *Phys. Rev. B (Rapid Commun.)* **36**, 822-825 (1987).

REFEREED PUBLICATIONS IN PRINT (cont'd)

95. "Antiferromagnetism of $\text{La}_2\text{CuO}_{4-y}$ Studied by Muon Spin Rotation," Y. J. Uemura, W. J. Kossler, X. H. Yu, J. R. Kempton, H. E. Schone, D. Opie, C. E. Stronach, D. C. Johnston, M. S. Alvarez, and D. P. Goshorn, *Phys. Rev. Lett.* **59**, 1045-1048 (1987).
96. "Soft X-ray Absorption Studies on High T_c Superconducting Oxides," D. Sondericker, Z. Fu, D. C. Johnston, and W. Eberhardt, *Phys. Rev. B (Rapid Commun.)* **36**, 3983-3985 (1987).
97. "Crystal Structure and Magnetic Properties of a New Molybdenophosphate: $\text{AgMo}_5\text{P}_8\text{O}_{33}$," K.-H. Lii, D. C. Johnston, D. P. Goshorn, and R. C. Haushalter, *J. Solid State Chem.* **71**, 131-138 (1987).
98. "Model for the Ground State of $\text{La}_{2z}\text{CuO}_{4-y}$," R. A. Klemm, H. Thomann, and D. C. Johnston, *Phys. Rev. B* **37**, 2239-2242 (1988).
99. "Neutron Diffraction Determination of Antiferromagnetic Structure of Cu Ions in $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ with $x = 0.0$ and 0.15 ," J. M. Tranquada, D. E. Cox, W. Kunmann, H. Moudden, G. Shirane, M. Suenaga, P. Zolliker, D. Vaknin, S. K. Sinha, M. S. Alvarez, A. J. Jacobson and D. C. Johnston, *Phys. Rev. Lett.* **60**, 156-159 (1988).
100. "Anomalous Ultrasound Propagation in High- T_c Superconductors: $\text{La}_{1.8}\text{Sr}_{0.2}\text{CuO}_{4-y}$ and $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$," S. Bhattacharya, M. J. Higgins, D. C. Johnston, A. J. Jacobson, J. P. Stokes, J. T. Lewandowsky and D. P. Goshorn, *Phys. Rev. B (Rapid Commun.)* **37**, 5901-5904 (1988).
101. "Systematic Variation of Magnetic Field Penetration Depth in High T_c Superconductors Studied by Muon Spin Relaxation," Y. J. Uemura, V. J. Emery, A. R. Moodenbaugh, M. Suenaga, D. C. Johnston, A. J. Jacobson, J. T. Lewandowski, J. H. Brewer, R. F. Kiefl, S. R. Kreitzman, G. M. Luke, T. Riseman, S. D. Dodds, C. E. Stronach, W. J. Kossler, J. R. Kempton, X. H. Yu, D. Opie and H. E. Schone, *Phys. Rev. B (Rapid Commun.)* **38**, 909-912 (1988).
102. "Elastic Anomalies and Phase Transitions in High- T_c Superconductors," S. Bhattacharya, M. J. Higgins, D. C. Johnston, A. J. Jacobson, J. P. Stokes, D. P. Goshorn and J. T. Lewandowski, *Phys. Rev. Lett.* **60**, 1181-1184 (1988).
103. "Structure and Magnetic Properties of $\text{V}_2(\text{VO})(\text{P}_2\text{O}_7)_2$. A Mixed Valence Vanadium (III,III,V) Pyrophosphate," J. W. Johnson, D. C. Johnston, H. E. King, Jr., T. R. Halbert, J. F. Brody and D. P. Goshorn, *Inorg. Chem.* **27**, 1646-1648 (1988).
104. "Superconductivity and Magnetism in the High T_c Copper Oxides," D. C. Johnston, S. K. Sinha, A. J. Jacobson and J. M. Newsam, *Physica C* **153-155**, 572-577 (1988).
105. Comparison between Muon Spin Rotation and Neutron Scattering Studies on the 3-Dimensional Magnetic Ordering of $\text{La}_2\text{CuO}_{4-y}$," Y. J. Uemura, W. J. Kossler, J. R. Kempton, X. H. Yu, H. E. Schone, D. Opie, C. E. Stronach, J. H. Brewer, R. F. Kiefl, S. R. Kreitzman, G. M. Luke, T. Riseman, D. L. Williams, E. J. Ansaldo, Y. Endoh, E. Kudo, K. Yamada, D. C. Johnston, M. Alvarez, D. P. Goshorn, Y. Hidaka, M. Oda, Y. Enomoto, M. Suzuki and T. Murakami, *Physica C* **153-155**, 769-770 (1988).

REFEREED PUBLICATIONS IN PRINT (cont'd)

106. "Antiferromagnetism in $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$," J. M. Tranquada, A. H. Moudden, A. I. Goldman, P. Zolliker, D. E. Cox, G. Shirane, S. K. Sinha, D. Vaknin, D. C. Johnston, M. S. Alvarez and A. J. Jacobson, *Phys. Rev. B* **38**, 2477-2485 (1988).
107. "Antiferromagnetism in the High- T_c Related Compounds," S. K. Sinha, D. E. Moncton, D. C. Johnston, D. Vaknin, G. Shirane and C. Stassis, *J. Appl. Phys.* **63**, 4015-4018 (1988).
108. "Ce Valence Mixing and Strong 3d Antiferromagnetism in CeMn_2Si_2 ," G. Liang, I. Perez, D. DiMarzio, M. Croft, D. C. Johnston, N. Anbalagan and T. Mihalison, *Phys. Rev. B (Rapid Commun.)* **37**, 5970-5973 (1988).
109. " $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4](\text{BF}_4)_2 \cdot \text{H}_2\text{O}$: Ferromagnetically Coupled Cr_4S Cluster with Spin 6 Ground State," A. Bino, D. C. Johnston, D. P. Goshorn, T. R. Halbert and E. I. Stiefel, *Science* **241**, 1479-1481 (1988).
110. "Observation of Triplet Hole Pairs and Glassy Spin Waves in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4-z}$ by Electron Spin Resonance," H. Thomann, R. A. Klemm, D. C. Johnston, P. J. Tindall, H. Jin, and D. P. Goshorn, *Phys. Rev. B* **38**, 6552-6560 (1988).
111. "Coexistence of Strong 3d Antiferromagnetism and Mixed-Valent Kondo Regime Transition in $\text{Ce}(\text{Mn}_x\text{Cr}_{1-x})_2\text{Si}_2$," G. Liang, M. Croft, D. C. Johnston, N. Anbalagan, and T. Mihalisin, *Phys. Rev. B* **38**, 5302-5309 (1988).
112. " μSR Studies on High- T_c Superconductivity," Y. J. Uemura, B. J. Sternlieb, D. E. Cox, V. J. Emery, A. Moodenbaugh, M. Suenaga, J. H. Brewer, J. F. Carolan, W. Hardy, R. Kadono, J. R. Kempton, R. F. Kiefl, S. R. Kreitzman, G. M. Luke, T. Riseman, D. L. Williams, B. X. Yang, W. J. Kossler, X. H. Yu, H. Schone, C. E. Stronach, J. Gopalakrishnan, M. A. Subramanian, A. W. Sleight, H. Hart, K. W. Lay, H. Takagi, S. Uchida, Y. Hidaka, T. Murakami, S. Etemad, D. Keane, V. Lee, J. B. Torrance, and D. C. Johnston, *J. Phys. (Paris) Colloque C8, Suppl.* **12**, 49, 2087-2092 (1988).
113. "Magnetic Susceptibility Study of a Model Supported Ziegler-Natta Catalyst: Evidence for Reduced Titanium Clusters," P. Brant, A. N. Specia and D. C. Johnston, *J. Catal.* **113**, 250-255 (1988).
114. "Synthesis of Organic Solids with a High Spin Density as an Approach to Organic Ferromagnets," L. Y. Chiang, D. C. Johnston, D. P. Goshorn, and A. N. Bloch, *Synthetic Metals* **27**, B639-B644 (1988).
115. "Synthesis and Properties of Non-stoichiometric Samples of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$; $0.04 \leq x \leq 1.00$," A. J. Jacobson, J. M. Newsam, D. C. Johnston, D. P. Goshorn, J. T. Lewandowski, and M. S. Alvarez, *Phys. Rev. B* **39**, 254-258 (1989).
116. "Superconductivity and Magnetism in the La_2CuO_4 -Based Compounds," Z. Fisk, S-W. Cheong and D. C. Johnston, *Mater. Res. Soc. Bull.* **14**, 33-36 (1989).

REFEREED PUBLICATIONS IN PRINT (cont'd)

117. "Mechanical and High Temperature (920°C) Magnetic Field Grain Alignment of Polycrystalline (Ho,Y)Ba₂Cu₃O_{7-δ}," A. Lusnikov, L. L. Miller, R. W. McCallum, S. Mitra, W. C. Lee, and D. C. Johnston, *J. Appl. Phys.* **65**, 3136-3141 (1989).
118. "Magnetic Susceptibility Scaling in La_{2-x}Sr_xCuO_{4-y}," D. C. Johnston, *Phys. Rev. Lett.* **62**, 957-960 (1989).
119. "Weakly Temperature-Dependent Highly Conducting Polyquinoline-Based Pyrolytic Films," L. Y. Chiang, J. P. Stokes, D. C. Johnston, and D. P. Goshorn, *Synthetic Metals* **29**, E483-E488 (1989).
120. "Synthesis of an Ambient Temperature Stable, High Spin Density Organic Solid with an Anomalously Small Interspin Coupling," L. Y. Chiang, D. C. Johnston, D. P. Goshorn, and A. N. Bloch, *J. Am. Chem. Soc. Commun.* **111**, 1925–1927 (1989).
121. "Superconducting Fluctuation Diamagnetism above T_c in YBa₂Cu₃O_{7-x}, La_{1.8}Sr_{0.2}CuO_{4-y} and Bi_{2-x}Pb_xSr₂CaCu₂O_{8+δ}," W. C. Lee, R. A. Klemm, and D. C. Johnston, *Phys. Rev. Lett.* **63**, 1012–1015 (1989).
122. "Antiferromagnetism in (Ca_{0.85}Sr_{0.15})CuO₂, the Parent of the Cuprate Family of Superconducting Compounds," D. Vaknin, E. Caignol, P. K. Davies, J. E. Fischer, D. C. Johnston and D. P. Goshorn, *Phys. Rev. B* **39**, 9122–9125 (1989).
123. "Magnetic Field Penetration Depth of Polycrystalline (Y,Gd)Ba₂Cu₃O₇, Grain-aligned YBa₂Cu₃O₇ and Single Crystal Bi₂Sr₂CaCu₂O₈," S. Mitra, J. H. Cho, W. C. Lee, D. C. Johnston, and V. G. Kogan, *Phys. Rev. B (Rapid Commun.)* **40**, 2674–2677 (1989).
124. "Neutron Scattering Studies of Antiferromagnetism in the High-T_c Compounds," S. K. Sinha, D. Vaknin, M. S. Alvarez, A. J. Jacobson, J. Newsam, J. T. Lewandowski, D. C. Johnston, C. Stassis, J. M. Tranquada, T. Freltoft, H. Moudden, A. I. Goldman, P. Zolliker, D. E. Cox and G. Shirane, *Physica B* **156 & 157**, 854–857 (1989).
125. "Magnetic Field Dependence of Ultrasound Velocity in High T_c Superconductors," M. J. Higgins, D. P. Goshorn, S. Bhattacharya, and D. C. Johnston, *Phys. Rev. B (Rapid Commun.)* **40**, 9393–9396 (1989).
126. "Low Temperature Behavior of Ni₃Al Alloys Near the Spin Fluctuator-Ferromagnet Phase Boundary," S. K. Dhar, K. A. Gschneidner, Jr., L. L. Miller, and D. C. Johnston, *Phys. Rev. B* **40**, 11488–11492 (1989).
127. "Synthesis, Structure, and Properties of Sr₂CuO₂Cl₂," L. L. Miller, X. L. Wang, S. X. Wang, C. Stassis, D. C. Johnston, J. Faber, Jr., and C.-K. Loong, *Phys. Rev. B* **41**, 1921–1925 (1990).
128. "Antiferromagnetism in Sr₂CuO₂Cl₂," D. Vaknin, S. K. Sinha, C. Stassis, L. L. Miller, and D. C. Johnston, *Phys. Rev. B* **41**, 1926–1933 (1990).
129. "Superconducting and Normal State Magnetic Susceptibility Anisotropy in YBa₂Cu₃O₇," W. C. Lee and D. C. Johnston, *Phys. Rev. B* **41**, 1904–1909 (1990).

REFEREED PUBLICATIONS IN PRINT (cont'd)

130. "Antiferromagnetic Form Factor of $\text{Sr}_2\text{CuO}_2\text{Cl}_2$," X. L. Wang, L. L. Miller, J. Ye, C. Stassis, B. N. Harmon, D. C. Johnston, A. J. Schultz, and C.-K. Loong, *J. Appl. Phys.* **67**, 4524–4526 (1990).
131. "Muon-Spin-Rotation Studies in Single Crystal $\text{Sr}_2\text{CuO}_2\text{Cl}_2$," L. P. Le, G. M. Luke, B. J. Sternlieb, Y. J. Uemura, J. H. Brewer, T. M. Riseman, D. C. Johnston, and L. L. Miller, *Phys. Rev. B* **42**, 2182–2187 (1990).
132. "New Phases Induced by Hydrogen Reduction and by Subsequent Oxidation of Ln_2CuO_4 (Ln = La, Pr, Nd, Sm, Eu, Gd)," F. C. Chou, J. H. Cho, L. L. Miller, and D. C. Johnston, *Phys. Rev. B* **42**, 6172–6180 (1990).
133. " μ +SR Studies on $\text{Sr}_2\text{CuO}_2\text{Cl}_2$, La_2CuO_4 , and Nd_2CuO_4 : 2-d Magnetism, Local Fields and Muon Sites," L. P. Le, G. M. Luke, B. J. Sternlieb, Y. J. Uemura, J. H. Brewer, T. M. Riseman, D. C. Johnston, L. L. Miller, Y. Hidaka, and H. Murakami, *Hyperfine Interactions* **63**, 279–286 (1990).
134. "Magnetic Susceptibility Anisotropy of Single Crystal $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$," D. C. Johnston and J. H. Cho, *Phys. Rev. B (Rapid Commun.)* **42**, 8710–8713 (1990).
135. "Magnetic Properties and Superconducting Fluctuation Diamagnetism above T_c in $\text{Bi}_{2-x}\text{Pb}_x\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ ($x = 0, 0.1, 0.2, 0.3, 0.5$) and $\text{Bi}_{2-x}\text{Pb}_x\text{Sr}_2\text{Ca}_2\text{Cu}_3\text{O}_{10+\delta}$ ($x = 0.2, 0.25$)," W. C. Lee, J. H. Cho, and D. C. Johnston, *Phys. Rev. B* **43**, 457–462 (1991).
136. "Heat Capacity of Single Crystal La_2CuO_4 and Polycrystalline $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ($0 \leq x \leq 0.20$) from 110 to 600 K," K. Sun, J. H. Cho, F. C. Chou, W. C. Lee, L. L. Miller, D. C. Johnston, Y. Hidaka, and T. Murakami, *Phys. Rev. B* **43**, 239–246 (1991).
137. "Heat Capacity of High Purity Polycrystalline $\text{YBa}_2\text{Cu}_3\text{O}_7$ from 0.4 K to 400 K in Applied Magnetic Fields of Zero and 70 kG," W. C. Lee, K. Sun, L. L. Miller, D. C. Johnston, R. A. Klemm, S. Kim, R. A. Fisher, and N. E. Phillips, *Phys. Rev. B* **43**, 463–472 (1991).
138. "The Antiferromagnetic Form Factor of La_2NiO_4 " X. L. Wang, C. Stassis, D. C. Johnston, T. C. Leung, J. Ye, B. N. Harmon, G. H. Lander, A. J. Schultz, C.-K. Loong, and J. M. Honig, *J. Appl. Phys.* **69**, 4860–4862 (1991).
139. "Magnetic and Electronic Properties of $\text{La}_x\text{Sr}_{1-x}\text{VO}_3$," A. V. Mahajan, D. C. Johnston, D. R. Torgeson, and F. Borsa, *Physica C* **185-189**, 1195–1196 (1991).
140. "Normal State Magnetism of the High T_c Cuprate Superconductors," D. C. Johnston, *J. Magn. Magn. Mater.* **100**, 218–240 (1991).
141. "A Scaling Property in the Reversible Magnetization of Anisotropic High- κ Type II Superconductors," Z. Hao, J. R. Clem, J. H. Cho, and D. C. Johnston, *Physica C* **185-189**, 1871–1872 (1991).
141. "Magnetic Penetration Depth of $\text{Ba}_{0.625}\text{K}_{0.375}\text{BiO}_3$," E. J. Ansaldo, Z. R. Wang, J. H. Cho, D. C. Johnston, and T. M. Riseman, *Physica C* **185-189**, 1889–1890 (1991).

REFEREED PUBLICATIONS IN PRINT (cont'd)

142. "Heat Capacity (150-300°K) and Anisotropic Magnetic Susceptibility (5-300°K) of Single Crystal $\text{La}_2\text{CuO}_{4+x}$," L. L. Miller, K. Sun, D. C. Johnston, J. E. Schirber, and Z. Fisk, *J Alloys Compds.* **183**, 312-324 (1992).
144. " ^{35}Cl NMR Study of Spin Dynamics in $\text{Sr}_2\text{CuO}_2\text{Cl}_2$," F. Borsa, M. Corti, T. Goto, A. Rigamonti, D. C. Johnston, and F. C. Chou, *Phys. Rev. B (Rapid Commun.)* **45**, 5756-5759 (1992).
145. "Neutron Diffraction Study of the Antiferromagnetic Form Factor of La_2NiO_4 ," X-L. Wang, C. Stassis, D. C. Johnston, T. C. Leung, J. Ye, B. N. Harmon, G. H. Lander, A. J. Schultz, C.-K. Loong, and J. M. Honig, *Phys. Rev. B* **45**, 5645-5653 (1992).
146. "Spin Dynamics in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ($0.02 \leq x \leq 0.08$) from ^{139}La NQR Relaxation: Fluctuations in a Finite Length Scale System," J. H. Cho, F. Borsa, D. C. Johnston, and D. R. Torgeson, *Phys. Rev. B (Rapid Commun.)* **46**, 3179-3182 (1992).
147. "Structural, Electronic and Magnetic Properties of $\text{La}_x\text{Sr}_{1-x}\text{VO}_3$ ($0.1 \leq x \leq 1.0$)", A. V. Mahajan, D. C. Johnston, D. R. Torgeson, and F. Borsa, *Phys. Rev. B* **46**, 10973-10985 (1992).
148. "Magnetic Properties of LaVO_3 ", A. V. Mahajan, D. C. Johnston, D. R. Torgeson, and F. Borsa, *Phys. Rev. B* **46**, 10966-10972 (1992).
149. "Magnetization Scaling of Single Crystal $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$ below T_c ", J. H. Cho, Zhidong Hao, and D. C. Johnston, *Phys. Rev. B (Rapid Commun.)* **46**, 8679-8682 (1992).
150. "Synthesis, Characterization, and Superconducting and Magnetic Properties of Electrochemically Oxidized $\text{La}_2\text{CuO}_{4+y}$ and $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4+y}$ ($0.01 \leq x \leq 0.33$, $0.01 \leq y \leq 0.36$)", F. C. Chou, J. H. Cho, and D. C. Johnston, *Physica C* **197**, 303-314 (1992).
151. "Phase Separation and Finite Size Scaling in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4+y}$ [$0 \leq (x,y) \leq 0.03$]", J. H. Cho, F. C. Chou, and D. C. Johnston, *Phys. Rev. Lett.* **70**, 222-225 (1993).
152. "Role of Vortex Fluctuations in Determining Superconducting Parameters from Magnetization Data for Layered Superconductors", V. G. Kogan, M. Ledvij, A. Yu. Simonov, J. H. Cho, and D. C. Johnston, *Phys. Rev. Lett.* **70**, 1870-1873 (1993).
153. "Structure of the Superconducting $\text{La}_2\text{CuO}_{4+y}$ Phases ($y \leq 0.08, 0.12$) Prepared by Electrochemical Oxidation", P. G. Radaelli, J. D. Jorgensen, A. J. Schultz, B. A. Hunter, J. L. Wagner, F. C. Chou, and D. C. Johnston, *Phys. Rev. B* **48**, 499-510 (1993).
154. "Magnetic Phase Diagram of Lightly Doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ from ^{139}La Nuclear Quadrupole Resonance", F. C. Chou, F. Borsa, J. H. Cho, D. C. Johnston, A. Lascialfari, D. R. Torgeson, and J. Ziolo, *Phys. Rev. Lett.* **71**, 2323-2326 (1993).
155. "Vortex Fluctuations in Polycrystalline $\text{Bi}_{1.75}\text{Pb}_{0.25}\text{Sr}_2\text{Ca}_2\text{Cu}_3\text{O}_{10}$ ", J. H. Cho, D. C. Johnston, M. Ledvij, and V. G. Kogan, *Physica C* **212**, 419-423 (1993).

REFEREED PUBLICATIONS IN PRINT (cont'd)

156. "Pressure Dependence of the Superconducting Transition Temperature of Superoxygenated $\text{La}_2\text{CuO}_{4+y}$ ", J. E. Schirber, W. E. Bayless, F. C. Chou, D. C. Johnston, P. C. Canfield, and Z. Fisk, *Phys. Rev. B* **48**, 6506-6508 (1993).
157. "Cu Spin Dynamics and Magnetic Quasiparticles in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$: a ^{139}La NQR Investigation", A. Lascialfari, F. Borsa, P. Ghigna, D. R. Torgeson, J. Ziolo, F. Chou, and D. C. Johnston, *J. Phys.: Condens. Matter* **5**, B19-B24 (1993).
158. "Preparation, Magnetization and Electrical Resistivity of Electrochemically Oxidized $\text{La}_2\text{CuO}_{4+x}$ Single Crystals", F. C. Chou, D. C. Johnston, S-W. Cheong, and P. C. Canfield, *Physica C* **216**, 66-76 (1993).
159. "Magnetic Penetration Depth in V_3Si and LiTi_2O_4 Measured by μSR ", W. D. Wu, A. Keren, L. P. Le, G. M. Luke, B. J. Sternlieb, Y. J. Uemura, D. C. Johnston, B. K. Cho, and P. Gehring, *Hyperfine Interactions* **86**, 615-621 (1994).
160. "Phase-Separation to Antiferromagnetic and Superconducting Domains in Oxygen Rich Single Crystal $\text{La}_2\text{CuO}_{4+x}$: Magnetic and Structural Neutron Diffraction Studies", D. Vaknin, J. L. Zarestky, D. C. Johnston, J. E. Schirber, and Z. Fisk, *Phys. Rev. B* **49**, 9057-9063 (1994).
161. "Phase Diagram and Phase Separation in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4+y}$ ", D. C. Johnston, F. Borsa, J. H. Cho, F. C. Chou, D. R. Torgeson, D. Vaknin, J. L. Zarestky, J. Ziolo, J. D. Jorgensen, P. G. Radaelli, A. J. Schultz, J. L. Wagner, and S.-W. Cheong, *J. Alloys Compds.* **207-208**, 206-212 (1994).
162. "Miscibility Gap in Electrochemically Oxygenated $\text{La}_2\text{CuO}_{4+y}$ ", P. G. Radaelli, J. D. Jorgensen, R. Kleb, B. A. Hunter, F. C. Chou and D. C. Johnston, *Phys. Rev. B* **49**, 6239-6245 (1994).
163. "Antiferromagnetic Ordering and Paramagnetic Behavior of Ferromagnetic Cu_6 and Cu_{18} Clusters in BaCuO_{2+x} ", Z.-R. Wang, X.-L. Wang, J. A. Fernandez-Baca, D. C. Johnston, and D. Vaknin, *Science* **264**, 402-404 (1994).
164. "Structural and Magnetic Studies of Sr_2IrO_4 ", M. K. Crawford, M. A. Subramanian, R. L. Harlow, J. A. Fernandez-Baca, Z. R. Wang, and D. C. Johnston, *Phys. Rev. B* **49**, 9198-9201 (1994).
165. "Superconducting State Magnetization and Critical Fields of Single Crystal $\text{YNi}_2\text{B}_2\text{C}$ ", M. Xu, P. C. Canfield, J. E. Ostenson, D. K. Finnemore, B. K. Cho, Z. R. Wang, and D. C. Johnston, *Physica C* **227**, 321-326 (1994).
166. "Magnetic Pair Breaking in $\text{HoNi}_2\text{B}_2\text{C}$ ", A. I. Goldman, C. Stassis, P. C. Canfield, J. Zarestky, P. Dervenagas, B. K. Cho, D. C. Johnston, and B. Sternlieb, *Phys. Rev. B (Rapid Commun.)* **50**, 9668-9671 (1994).
167. "Phase Separation and Doped-Hole Segregation in $\text{La}_2\text{CuO}_{4+y}$ and $\text{La}_{2-x}\text{Sr}_x\text{CuO}_{4+y}$ ", D. C. Johnston, W. R. Bayless, F. Borsa, P. C. Canfield, S-W. Cheong, J. H. Cho, F. C. Chou, Z. Fisk, J. D. Jorgensen, A. Lascialfari, L. L. Miller, P. G. Radaelli, J. E. Schirber, A. J. Schultz, D. R. Torgeson, D. Vaknin, J. L. Wagner, J. Zarestky, and J. Ziolo, *Physica C* **235-240**, 257-260 (1994).

REFEREED PUBLICATIONS IN PRINT (cont'd)

168. "Specific Heat and Anisotropic Superconducting and Normal-State Magnetization of $\text{HoNi}_2\text{B}_2\text{C}$ ", P. C. Canfield, B. K. Cho, D. C. Johnston, D. K. Finnemore, and M. F. Hundley, *Physica C* **230**, 397-406 (1994).
169. "Magnetization, Upper Critical Field, and Isotropy in Single Crystals $\text{YNi}_2\text{B}_2\text{C}$ ", M. Xu, B. K. Cho, P. C. Canfield, D. K. Finnemore, and D. C. Johnston, *Physica C* **235-240**, 2533-2534 (1994).
170. "Oxygen Diffusion in $\text{La}_2\text{CuO}_{4+y}$ from ^{139}La NMR-NQR Relaxation," S. Rubini, F. Borsa, P. Canfield, F. C. Chou, Q. Hu, A. Lascialfari, D. C. Johnston, A. Rigamonti, and D. R. Torgeson, *Physica C* **235-240**, 1717-1718 (1994).
171. "Hole Segregation and Spin and Charge Localization in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ from ^{139}La NQR and $\mu\text{+SR}$ ", F. Borsa, P. Carretta, A. Lascialfari, D. R. Torgeson, F. C. Chou, and D. C. Johnston, *Physica C* **235-240**, 1713-1714 (1994).
172. " Sr_2IrO_4 and Sr_2RhO_4 : Structural and Magnetic Studies of 4d and 5d Transition Metal Analogs of La_2CuO_4 ," M. A. Subramanian, M. K. Crawford, R. L. Harlow, T. Ami, J. A. Fernandez-Baca, Z. R. Wang, and D. C. Johnston, *Physica C* **235-240**, 743-744 (1994).
173. "Magnetic Susceptibility and Low-Temperature Structure of the Linear Chain Cuprate Sr_2CuO_3 ", T. Ami, M. K. Crawford, R. L. Harlow, Z. R. Wang, D. C. Johnston, Q. Huang, and R. W. Erwin, *Phys. Rev. B* **51**, 5994-6001 (1995).
174. "Influence of Temperature and Magnetic Field on Ferrimagnetic CePtGe_2 ", V. K. Pecharsky, L. L. Miller, K. A. Gschneidner, Jr., and D. C. Johnston, *Phys. Rev. B* **51**, 954-961 (1995).
175. "Magnetic Structure of $\text{ErNi}_2\text{B}_2\text{C}$ ", J. Zarestky, C. Stassis, A. I. Goldman, P. C. Canfield, P. Dervenagas, B. K. Cho, and D. C. Johnston, *Phys. Rev. B (Rapid Commun.)* **51**, 678-680 (1995).
176. "Neutron Diffraction Study of the Magnetic Ordering of BaCuO_{2+x} ", Xun-Li Wang, J. A. Fernandez-Baca, Z. R. Wang, D. Vaknin, and D. C. Johnston, *Physica B* **213-214**, 97-99 (1995).
177. "Magnetic and Superconducting Properties of Single Crystal $\text{TmNi}_2\text{B}_2\text{C}$ ", B. K. Cho, M. Xu, P. C. Canfield, L. L. Miller, and D. C. Johnston, *Phys. Rev. B* **52**, 3676-3683 (1995).
178. "Magnetism and Superconductivity in Single Crystal $\text{ErNi}_2\text{B}_2\text{C}$ ", B. K. Cho, P. C. Canfield, L. L. Miller, D. C. Johnston, W. P. Beyermann, and A. Yatskar, *Phys. Rev. B* **52**, 3684-3695 (1995).
179. "Onset of Superconductivity in the Antiferromagnetically Ordered State of Single Crystal $\text{DyNi}_2\text{B}_2\text{C}$ ", B. K. Cho, P. C. Canfield and D. C. Johnston, *Phys. Rev. B (Rapid Commun.)* **52**, R3844-R3847 (1995).

REFEREED PUBLICATIONS IN PRINT (cont'd)

180. "Superconducting Anisotropy of $\text{YNi}_2\text{B}_2\text{C}$ ", E. Johnston-Halperin, J. Fiedler, D. E. Farrell, M. Xu, B. K. Cho, P. C. Canfield, D. K. Finnemore, and D. C. Johnston, *Phys. Rev. B* **51**, 12852-12853 (1995).
181. "Spin Correlations and Magnetic Field Effects in the Weakly Anisotropic Square-Lattice Antiferromagnet $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ ", B. J. Suh, F. Borsa, L. L. Miller, M. Corti, D. C. Johnston, and D. R. Torgeson, *Phys. Rev. Lett.* **75**, 2212-2215 (1995); 75, 4335 (E) (1995).
182. "Staggered Magnetization in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ from ^{139}La NQR and μSR : Effects of Sr Doping in the Range $0 < x < 0.02$ ", F. Borsa, P. Carretta, J. H. Cho, F. C. Chou, Q. Hu, D. C. Johnston, A. Lascialfari, D. R. Torgeson, R. J. Gooding, N. M. Salem, and K. J. E. Vos, *Phys. Rev. B* **52**, 7334-7345 (1995).
183. "Energy Gap of Quaternary Compound Superconductor $\text{YNi}_2\text{B}_2\text{C}$ and $\text{LuNi}_2\text{B}_2\text{C}$ by Point Contact Spectroscopy", G. T. Jeong, J. I. Kye, S. H. Chun, Z. G. Khim, W. C. Lee, P. C. Canfield, B. K. Cho, and D. C. Johnston, *Physica C* **253**, 48-52 (1995).
184. "Static Magnetization and AC Susceptibility Measurements of the Copper-Oxygen Cluster Compound BaCuO_{2+x} ", Z. R. Wang, D. C. Johnston, L. L. Miller, and D. Vaknin, *Phys. Rev. B* **52**, 7384-7394 (1995).
185. "Superconductivity in the Transition Metal-Rich Layered Compound $\text{Ta}_x\text{Nb}_{5-x}\text{S}_2$ ($x = 1.72$)", Zhaorong Wang, David C. Johnston, Xiaoqiang Yao, and Hugo F. Franzen, *Physica C* **252**, 138-140 (1995).
186. "Optical Properties of Single Crystal $\text{La}_2\text{CuO}_{4+y}$ ", M. A. Quijada, D. B. Tanner, F. C. Chou, D. C. Johnston, and S-W. Cheong, *Phys. Rev. B* **52**, 15485-15503 (1995).
187. "Oxygen Ordering and Phase Separation in $\text{La}_2\text{CuO}_{4+y}$ ", B. W. Statt, P. C. Hammel, Z. Fisk, S-W. Cheong, F. C. Chou, D. C. Johnston, and J. E. Schirber, *Phys. Rev. B* **52**, 15575-15581 (1995).
188. "Superconducting and Normal State Magnetic Properties of $\text{RNi}_2\text{B}_2\text{C}$ Single Crystals", D. C. Johnston, F. Borsa, B. K. Cho, P. C. Canfield, P. Dervenagas, A. I. Goldman, B. N. Harmon, L. L. Miller, C. Stassis, B. J. Suh, D. R. Torgeson, M. Xu, J. Zarestky, M. F. Hundley, R. Movshovich, J. D. Thompson, A. V. Chubukov, and B. Sternlieb, *Chinese J. Phys.* **34**, 397-405 (1996).
189. "Crystalline Electric Field Effects in Single Crystal $\text{HoNi}_2\text{B}_2\text{C}$ ", B. K. Cho, B. N. Harmon, D. C. Johnston, and P. C. Canfield, *Phys. Rev. B* **53**, 2217-2220 (1996).
190. "Intercalation and Staging Behavior in Super-Oxygenated $\text{La}_2\text{CuO}_{4+y}$ " B. O. Wells, R. J. Birgeneau, F. C. Chou, Y. Endoh, D. C. Johnston, M. A. Kastner, Y. S. Lee, G. Shirane, J. M. Tranquada, and K. Yamada, *Z. Phys. B* **100**, 535-545 (1996).
191. "Evidence for Crossover Effects in the Spin Dynamics of the Two-Dimensional Antiferromagnet $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ from ^{35}Cl Nuclear Magnetic Resonance", B. J. Suh, F. Borsa, L. L. Miller, D. C. Johnston, D. R. Torgeson, and M. Corti, *J. Appl. Phys.* **79**, 5084-5086 (1996).

REFEREED PUBLICATIONS IN PRINT (cont'd)

192. "Phase Separation Kinetics in $\text{La}_2\text{CuO}_{4+y}$ and Inhomogenous Hole Doping in the Antiferromagnetic Regime ($0 < x < 0.02$) of $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ", D. C. Johnston, F. C. Chou, F. Borsa, P. Carretta, A. Lascialfari, R. J. Gooding, N. M. Salem, K. J. E. Vos, J. H. Cho, and D. R. Torgeson, *J. Supercond.* **9**, 337-342 (1996).
193. "Absence of Antiferromagnetic Correlations in $\text{YNi}_2\text{B}_2\text{C}$ ", B. J. Suh, F. Borsa, D. R. Torgeson, B. K. Cho, P. C. Canfield, D. C. Johnston, J. Y. Rhee, and B. N. Harmon, *Phys. Rev. B (Rapid Commun.)* **53**, R6022-R6025 (1996).
194. "Phase Separation and Oxygen Diffusion in Electrochemically Oxidized $\text{La}_2\text{CuO}_{4+y}$: A Static Magnetic Susceptibility Study", F. C. Chou and D. C. Johnston, *Phys. Rev. B* **54**, 572-583 (1996).
195. "Magnetic Anisotropy and Weak Ferromagnetism of Single Crystal $\text{TbNi}_2\text{B}_2\text{C}$ ", B. K. Cho, P. C. Canfield, and D. C. Johnston, *Phys. Rev. B* **53**, 8499-8505 (1996).
196. "Vortex Glass and Lattice Melting Transitions in a $\text{YNi}_2\text{B}_2\text{C}$ Single Crystal", Mi-Ock Mun, Sung-Ik Lee, W. C. Lee, P. C. Canfield, B. K. Cho, and D. C. Johnston, *Phys. Rev. Lett.* **76**, 2790-2793 (1996).
197. "Raman Modes of $\text{RNi}_2\text{B}_2\text{C}$ ($\text{R} = \text{Lu}, \text{Ho}, \text{Y}$) Single Crystals", Hyun-Jung Park, Hye-Soo Shin, Hye-Gyong Lee, In-Sang Yang, W. C. Lee, B. K. Cho, P. C. Canfield, and D. C. Johnston, *Phys. Rev. B* **53**, 2237-2240 (1996).
198. "Antiferromagnetic Exchange in Two-Leg Spin-1/2 Ladders", D. C. Johnston, *Phys. Rev. B* **54**, 13009-13016 (1996).
199. "Breakdown of de Gennes Scaling in $(\text{R}_{1-x}\text{R}_x)\text{Ni}_2\text{B}_2\text{C}$ Compounds", B. K. Cho, P. C. Canfield, and D. C. Johnston, *Phys. Rev. Lett.* **77**, 163-166 (1996).
200. "Nonlocal Electrodynamics and Low -Temperature Magnetization of Clean High-Superconductors", V. G. Kogan, A. Gurevich, J. H. Cho, D. C. Johnston, M. Xu, J. R. Thompson, and A. Martynovich, *Phys. Rev. B* **54**, 12386-12396 (1996).
201. "Electronic and Magnetic Properties of Single Crystal $\text{YNi}_2\text{B}_2\text{C}$ from ^{11}B and ^{89}Y NMR and Magnetic Susceptibility Measurements", B. J. Suh, F. Borsa, D. R. Torgeson, B. K. Cho, P. C. Canfield, D. C. Johnston, J. Y. Rhee, and B. N. Harmon, *Phys. Rev. B* **54**, 15341-15351 (1996).
202. "Tunneling Measurement for the Energy Gap in Quaternary Compound Superconductors $\text{YNi}_2\text{B}_2\text{C}$, $\text{LuNi}_2\text{B}_2\text{C}$ and YBCO by LTSTM", G. T. Jeong, J. I. Kye, S. H. Chun, W. K. Park, Z. G. Khim, W. C. Lee, P. C. Canfield, B. K. Cho, and D. C. Johnston, *Chinese J. Phys.* **34**, 406-413 (1996).
203. "Magnetic Susceptibility of Sr_2CuO_3 and SrCu_2O_3 ", D. C. Johnston, *Acta Phys. Pol. A* **91**, 181-189 (1997).

REFEREED PUBLICATIONS IN PRINT (cont'd)

204. "Elastic and Quasielastic Neutron Scattering Study of the Spin-1/2 Square Lattice Antiferromagnet $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ ", D. Vaknin, L. L. Miller, J. L. Zarestky, and D. C. Johnston, *Physica C* **274**, 331-341 (1997).
205. "LiV₂O₄: A Heavy Fermion Transition Metal Oxide", S. Kondo, D. C. Johnston, C. A. Swenson, F. Borsa, A. V. Mahajan, L. L. Miller, T. Gu, A. I. Goldman, M. B. Maple, D. A. Gajewski, E. J. Freeman, N. R. Dilley, R. P. Dickey, J. Merrin, K. Kojima, G. M. Luke, Y. J. Uemura, O. Chmaissem, and J. D. Jorgensen, *Phys. Rev. Lett.* **78**, 3729-3732 (1997).
206. "Temperature Dependent Modulation Amplitude of the CuO₂ Superconducting Lattice in La₂CuO_{4.1}", A. Lanzara, N. L. Saini, A. Bianconi, J. L. Hazemann, Y. Soldo, F. C. Chou, and D. C. Johnston, *Phys. Rev. B* **55**, 9120-9124 (1997).
207. "Structure and Thermal Expansion of LiV₂O₄: Correlation Between Structure and Heavy Fermion Behavior", O. Chmaissem, J. D. Jorgensen, S. Kondo, and D. C. Johnston, *Phys. Rev. Lett.* **79**, 4866-4869 (1997).
208. "Localized Holes in Superconducting Lanthanum Cuprate", P. C. Hammel, B. W. Statt, R. L. Martin, F. C. Chou, D. C. Johnston, and S-W. Cheong, *Phys. Rev. B (Rapid Commun.)* **57**, R712-R715 (1998).
209. "Magnetic Neutron Diffraction from the Magnetic Clusters in Single Crystal BaCuO_{2+x}", D. Vaknin, J. P. Koster, J. L. Zarestky, Z. R. Wang, J. A. Fernandez-Baca, and D. C. Johnston, *Phys. Rev. B* **57**, 465-470 (1998).
210. "⁷Li and ⁵¹V NMR Study of the Heavy Fermion Compound LiV₂O₄", A. V. Mahajan, R. Sala, E. Lee, F. Borsa, S. Kondo, and D. C. Johnston, *Phys. Rev. B* **57**, 8890-8899 (1998).
211. "Muon Spin Relaxation Measurements in LiV₂O₄", J. Merrin, Y. Fudamoto, K. M. Kojima, M. Larkin, G. M. Luke, B. Nachimi, Y. J. Uemura, S. Kondo, and D. C. Johnston, *J. Magn. Magn. Mater.* **177-181**, 799-800 (1998).
212. "Superfluid and Normal Fluid Density in High-T_c Superconductors," D. B. Tanner, H. L. Liu, M. A. Quijada, A. M. Zibold, H. Berger, R. J. Kelley, M. Onellion, F. C. Chou, D. C. Johnston, J. P. Rice, D. M. Ginsberg, and J. T. Markert, *Physica B* **244**, 1-8 (1998).
213. "Study of Cu-O and Cu-La Pair Correlations in Superconducting La₂CuO_{4.1} System by Cu K-Edge EXAFS," N. L. Saini, A. Lanzara, A. Bianconi, F. C. Chou, and D. C. Johnston, *J. Phys. Soc. Jpn.* **67**, 16-19 (1998).
214. "Quantum Monte Carlo Simulation of the Trellis Lattice Heisenberg Model for SrCu₂O₃ and CaV₂O₅," S. Miyahara, M. Troyer, D. C. Johnston, and K. Ueda, *J. Phys. Soc. Jpn.* **67**, 3918-3923 (1998).
215. "Anomalous Structural Modulations in La₂CuO_{4.1}", A. Lanzara, N. L. Saini, A. Bianconi, F. C. Chou, and D. C. Johnston, *Physica C* **296**, 1-2 (1998).

REFEREED PUBLICATIONS IN PRINT (cont'd)

216. "Synthesis, Characterization and Magnetic Susceptibility of the Heavy Fermion Transition Metal Oxide LiV_2O_4 ", S. Kondo, D. C. Johnston, and L. L. Miller, *Phys. Rev. B* **59**, 2609-2626 (1999).
217. "Specific Heat and Thermal Expansion of the Heavy Fermion Compound LiV_2O_4 ", D. C. Johnston, C. A. Swenson, and S. Kondo, *Phys. Rev. B* **59**, 2627-2641 (1999).
218. "Spin Dynamics in Hole-Doped Two-Dimensional $S = 1/2$ Heisenberg Antiferromagnets: ^{63}Cu NQR Relaxation in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ for $x \leq 0.04$ ", P. Carretta, F. Tedoldi, A. Rigamonte, F. Galli, F. Borsa, J. H. Cho, and D. C. Johnston, *Eur. Phys. J. B* **10**, 233-236 (1999).
219. "The Low-Temperature Specific Heat of BaCuO_2 and $\text{BaCuO}_{2.14}$ in Magnetic Fields to 7 T", R. A. Fisher, D. A. Wright, J. P. Emerson, B. F. Woodfield, N. E. Phillips, Z.-R. Wang, and D. C. Johnston, *Phys. Rev. B* **61**, 538-548 (2000).
220. "Polynuclear Chromium(III) Carboxylates. 1. Synthesis, Structure and Magnetic Properties of an Octanuclear Complex with a Ring Structure", M. Eshel, A. Bino, I. Felner, D. C. Johnston, M. Luban, and L. L. Miller, *Inorg. Chem.* **39**, 1376-1380 (2000).
221. "Magnetism and Spin Dynamics in the Cluster Compound $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4](\text{NO}_3)_2 \cdot \text{H}_2\text{O}$ ", Y. Furukawa, M. Luban, F. Borsa, D. C. Johnston, A. V. Mahajan, L. L. Miller, D. Mentrup, J. Schnack, and A. Bino, *Phys. Rev. B* **61**, 8635-8638 (2000).
222. "Thermodynamics of Spin $S = 1/2$ Antiferromagnetic Uniform and Alternating-Exchange Heisenberg Chains", D. C. Johnston, R. K. Kremer, M. Troyer, X. Wang, A. Klümper, S. L. Bud'ko, A. F. Panchula, and P. C. Canfield, *Phys. Rev. B* **61**, 9558-9606 (2000).
223. "Thermodynamics of the Spin-1/2 Antiferromagnetic Uniform Heisenberg Chain", A. Klümper and D. C. Johnston, *Phys. Rev. Lett.* **84**, 4701-4704 (2000).
224. "Heavy Fermion Behaviors in LiV_2O_4 ", D. C. Johnston, *Physica B* **281-282**, 21-25 (2000); invited review paper.
225. "Single Crystal Growth of the High Pressure Phase of $(\text{VO})_2\text{P}_2\text{O}_7$ at 3 GPa", T. Saito, T. Terashima, M. Azuma, M. Takano, T. Goto, H. Ohta, W. Utsumi, P. Bordet, and D. C. Johnston, *J. Solid State Chem.* **153**, 124-131 (2000).
226. "Coexistence of Stripes and Superconductivity: Tc Amplification in a Superlattice of Superconducting Stripes", A. Bianconi, D. DiCastro, G. Bianconi, A. Pifferi, N. L. Saini, F. C. Chou, D. C. Johnston, and M. Colapietro, *Physica C* **341**, 1719-1722 (2000).
227. "Modeling of the Magnetic Susceptibilities of the Ambient- and High-Pressure Phases of $(\text{VO})_2\text{P}_2\text{O}_7$ ", D. C. Johnston, T. Saito, M. Azuma, M. Takano, T. Yamauchi, and Y. Ueda, *Phys. Rev. B* **64**, 134403 (2001), 23 pages.
228. "Decomposition of the Spin-1/2 Heisenberg Chain Compound Sr_2CuO_3 in Air and Water: an EPR and Magnetic Susceptibility Study of $\text{Sr}_2\text{Cu}(\text{OH})_6$ ", J. M. Hill, D. C. Johnston, and L. L. Miller, *Phys. Rev. B* **65**, 134428 (2002), 9 pages.

REFEREED PUBLICATIONS IN PRINT (cont'd)

229. “Magnetization Study of the Ultrahard Material MgAlB_{14} ,” J. M. Hill, D. C. Johnston, B. A. Cook, J. L. Harringa, and A. M. Russell, *J. Magn. Magn. Mater.* (accepted).
230. “Examining the Metal-to-Insulator Transitions in $\text{Li}_{1+x}\text{Ti}_{2-x}\text{O}_4$ and $\text{LiAl}_y\text{Ti}_{2-y}\text{O}_4$ with a Quantum Site Percolation Model”, F. Fazileh, R. J. Gooding, and D. C. Johnston, *Phys. Rev. B* 69, 104503 (2004), 5 pages.
- 230A. “Magnetic Susceptibilities of Spin-1/2 Antiferromagnetic Heisenberg Ladders and Applications to Ladder Oxide Compounds”, D. C. Johnston, M. Troyer, S. Miyahara, D. Lidsky, K. Ueda, M. Azuma, Z. Hiroi, M. Takano, M. Isobe, Y. Ueda, M. A. Korotin, V. I. Anisimov, A. V. Mahajan, and L. L. Miller, *Phys. Rev. B* (submitted), 63 typeset pages with 56 figures.